

CONTENTS

	Page
Biology of <i>Caryedon crineus</i> Arora (Coleoptera: Bruchidae), a seed pest of <i>Cassia occidentalis</i> (L.): D. R. Thakur, H. S. Banyal.	1
Ecdysone-neurosecretion interaction during vitellogenesis in <i>Poecilocerus pictus</i> Fabr. (Orthoptera: Acrididae): Versha Sharma.	9
Study of beetles (Coleoptera) associated with stored products in Kolkata, India 1. Genus <i>Carpophilus</i> Stephens (Nitidulidae): P. K. Basak, T. K. Pal.	17
Impact of IPM on natural enemies in irrigated cotton of North India: R. K. Tanwar, O. M. Bambawale, P. Jeyakumar, A. Dhandapani, Vikas Kanwar, O. P. Sharma, D. Monga.	25
Super parasitism in Indian lac insect, <i>Kerria lacca</i> (Kerr) and its implication on fecundity and resin producing efficiency of its two strains: K. K. Sharma, Kavita Kumari, Suman Lakhanpaul.	33
New species of <i>Thelcticopis</i> (Araneae: Sparassidae) from India: K. Sunil Jose, P. A. Sebastian.	41
Activity pattern of juvenile hormone during metamorphosis in <i>Dysdercus cingulatus</i> Fabr. (Heteroptera: Pyrrhocoridae): U. Gayathri Elayidam, D. Muraleedharan.	47
SHORT COMMUNICATIONS	
Assessment of storage losses in wheat due to stored grain insect pests in Himachal Pradesh: K. R. S. Thakur, P. K. Mehta, Meena Thakur.	53
The suitability of host plants for mass rearing of <i>Scirtothrips dorsalis</i> Hood (Thysanoptera: Thripidae): P. N. Ganga Visalakshy, A. Krishnamoorthy.	57
Ant fauna associated with whiteflies in some locales of Eastern and Western Ghats in two Southern states of India: R. W. Alexander Jesudasan, M. Anto Claver.	61
Histochemical study of yolk granules of ripe oocytes of <i>Oxya hyla hyla</i> (Orthoptera: Acrididae): K. Das, D. Ghosh.	67



AUTHOR INDEX

Alexander Jesudasan, R. W., 61	Lakhanpaul, Suman , 33
Anto Claver, M., 61	
Bambawale, O. M., 25	Mehta, P. K., 53
Banyal, H. S., 1	Monga, D., 25
Basak, P. K., 17	Muraleedharan, D., 47
Das, K., 67	
Dhandapani, A., 25	Pal, T. K., 17
Ganga Visalakshy, P. N., 57	Sebastian, P. A., 41
Gayathri Elayidam, U., 47	Sharma, O. P., 25
Ghosh, D., 67	Sharma, Versha , 9
Jeyakumar, P., 25	Sharma, K. K., 33
Kanwar, Vikas , 25	Sunil Jose, K., 41
Krishnamoorthy, A., 57	
Kumari, Kavita , 33	Tanwar, R. K., 25
	Thakur, K. R. S., 53
	Thakur , Meena , 53
	Thakur, D. R., 1

Statement of ownership and other particulars about ENTOMON

(Form IV, Rule 8 of Registration of Newspapers (Central) Rules 1956)

1. Place of publication:	Trivandrum
2. Periodicity of publication:	Quarterly
3. Printer's name, nationality: and address:	D. Muraleedharan, Indian CABB, University of Kerala Kariavattom, Trivandrum 695581
4. Publisher's name, nationality and address:	-do-
5. Editor's name, nationality and address:	-do-
6. Name and address of the individual who owns the newspaper:	Association for Advancement of Entomology Department of Zoology, University of Kerala Kariavattom, Trivandrum 695581

I, D. Muraleedharan, hereby declare that the particulars given above are true to the best of my knowledge and belief.

Trivandrum
31 March 2007

Sd/-
D. Muraleedharan
Publisher, ENTOMON



Biology of *Caryedon crineus* Arora (Coleoptera: Bruchidae), a seed pest of *Cassia occidentalis* (L.)

D. R. Thakur* and H. S. Banyal

Department of Biosciences, Himachal Pradesh University, Shimla 171005, India
Email: drdr4@rediffmail.com

ABSTRACT: Biology and behaviour of *Caryedon crineus* Arora, infesting pods and seeds of *Cassia occidentalis* (L.) was studied. Female beetle lays 16–27 eggs on pods and seeds of host plant during nights, which hatch after about 9 days. All larval instars are voracious feeders and their development is completed inside the host seed. Larval duration lasts for 23–27 days and last instar larva emerges out of seed by cutting a hole and then undergoes pupation outside. Pupal duration inside cocoon lasts for 51–60 days during favourable period and 191–220 days during adverse conditions and adult emerged out after 85–96 days. © 2007 Association for Advancement of Entomology

KEYWORDS: *Caryedon crineus*, biology, *Cassia occidentalis*

INTRODUCTION

Caryedon crineus Arora (Bruchidae: Coleoptera) is a pest of *Cassia occidentalis* (L.) (Panwar) in India (Arora, 1977). Adults appear in the field in the month of April after passing the winter as hibernated pupa. Biology and ecology of species of *Caryedon* have been studied by various workers. Pajni and Mann (1979); Pierre and Huignard (1990) studied the biology of *Caryedon serratus* (Oliver). Conway (1983) showed that mating, oviposition and development in *C. serratus* occurred in considerable depth in jute bags but fourth instar pupates outside the bag stacks. Halle *et al.* (2002) studied the comparative biology of tamarind beetle, *C. serratus* on nine different hosts in laboratory and Thakur and Banyal (2003) demonstrated the life history of *C. crineus*. Southgate (1979) observed three different sites of cocoon formation among some Nigerian species of genus *Caryedon*. The taxonomic position of *C. crineus* was reported by Arora (1977, 1978). Except for life history and larval parasitoid (Thakur and Banyal, 2003, 2005) there is no information on biological aspects of this species. Hence studies were made on the biology and behaviour of *C. crineus*.

*Corresponding author

MATERIALS AND METHODS

Adults of *C. crineus* emerged in the month of April were kept in pairs (one male and one female) along with seeds and pods of *C. occidentalis* in wire mesh cages of $10 \times 12 \times 12 \text{ cm}^3$ and 500 ml glass jars with their open end covered with muslin cloth, in the laboratory. Observations were made on behaviour of adults and immature stages. The number of eggs laid per female, their shape and size and incubation period were also observed. Morphometric studies of eggs and first instar larvae were carried out. Eggs of almost equal age were marked and observations were made daily. The larval instars duration, pupal duration, time taken for completion of life cycle, longevity of insects and number of generations completed in a year were recorded.

RESULTS

Behaviour

The male *Caryedon crineus* Arora was smaller in size than the female and had vertical pygidium. Adults of both the sexes showed brisk activity during the evening and early morning hours and remained hidden and inactive during daytime. Adults feigned death when touched or disturbed, turned upside down, retracted neck, pulled their legs and antennae close to the body and remained motionless but regained normal posture and activities after 10–20 seconds. Mating occurred inside the pods among the seeds during night time because neither mating nor egg laying has been observed during day time.

Mating

Male recognized the female insect from a close distance, chased her and continuously tapped her back or pygidium with its antennae. Male then put its antennae on the back of the female and the female remained motionless. Male insect then mounted on the back of the female and held her abdomen with his fore and mid legs. During the initial time of mating, the female pushed the male off her back with her hind legs, but persistent attempts of the male facilitated the insertion of already extruded aedeagus into the genital opening of the female resulting in successful copulation. During copulation female remained stationary and jerking movement of the male insect was observed. When another male tried to dislodge the copulating couple, the female moved ahead and pulled the inseminating male on her back.

After 10–20 min of copulation the female pushed the male away with her hind legs. After copulation male and female cleaned their mouthparts and antennae with fore and mid legs and genitalia or ovipositor with hind legs and then started moving away. When two males tried to mount the same female, the stronger one succeeded in the attempt and weaker moved away in search of another receptive female.

Oviposition

Gravid female of *C. crineus* started laying eggs 7.1 ± 1.1 days after emergence and eggs were laid during night time only. Females of *C. crineus* followed one seed -

one egg pattern but pods of the host plant contained 1–8 eggs. Females of *C. crineus* after laying their full quota of eggs became sluggish and died 3.3 ± 1.3 days after the oviposition was completed.

Before depositing an egg, the female examined the seed carefully with her antennae. It even turned over the seeds by thrusting her head underneath the seeds and pushed them. After searching a seed, the female lowered her abdomen, extruded the ovipositor, rubbed it on the surface of the seed and released a sticky fluid for attachment of the egg. The egg was then deposited on the treated surface with its broader end coming out first from the ovipositor by shaking abdomen. The whole process of depositing an egg on the surface of seed or pod took about 2–3 min. Oviposition period of a female of *C. crineus* extended for 9.0 ± 2.2 days and maximum of 9 eggs were deposited on the 2nd night of her oviposition period.

A single female of *C. crineus* laid 21.8 ± 3.2 eggs. Freshly laid eggs were rusty or light orange, oval in shape and measured 0.84 ± 0.06 mm in length and 0.496 ± 0.03 mm in width. Eggs laid in the beginning and near the end of oviposition period were empty and without ooplasm and failed to hatch. In the absence of seeds egg laying was postponed and in the prolonged absence of the seeds eggs were deposited on the walls of the test tube or petridish. Such eggs on the walls of culture vessels produced normal first instar larvae inside the egg shell but the larvae did not hatch out and died inside the egg shell within 2–3 days.

Incubation period lasted for 8.9 ± 0.37 days under natural atmospheric conditions (24°C – 39°C temperature and 53%–98% relative humidity). Egg shell became transparent before hatching and developing first instar larvae were visible. But this visibility was lost before hatching due to deposition of the larval frass inside the egg shells.

Larval development

First instar larvae hatched out from the ventral surface of the eggs, penetrated vertically into the host seeds and then turned at an angle to feed on the contents of the host seeds. On the way into the host seed first instar larva consumed the contents of seed and whitish and light orange frass was deposited inside the empty egg shell. The entrance hole into the seed was closed by the first instar larva with its faecal matter but could be identified as a white mark on the seed.

First instar larvae measured 0.469 ± 0.02 mm long and 0.237 ± 0.01 mm broad, each having large brownish rounded head and elongated tapering body covered with numerous white shining setae of variable size. The body of larva was divided into head, thorax and abdomen. Head possessed the necessary appendages, viz. clypeus, labrum, labium, a pair of mandibles, a pair of maxillae and a pair of antennae. Thorax possessed three pairs of legs and abdomen was tapering posteriorly and devoid of any appendages. First instar larva possessed H- shaped prothoracic plate with six pairs of dark, highly sclerotized teeth at its outer margin. The prothoracic plate facilitated the entry of larva into the host seed. Second and successive instar larvae did not possess the prothoracic plate. Each larval instar exuded a yellowish fluid to soften the contents of the seed. Head was smaller but deeply retracted in prothorax in successive instar

larvae. All the larval instars were voracious feeders and kept on feeding all the time except during moulting and consumed the seed contents completely leaving behind seed coat only.

Further development of first and successive instar larvae was completed inside the host seed. Larvae did not shift from one seed to another even when the contents of the initially infested seed were exhausted probably due to the inability of second and successive instars to possess the chiseling apparatus, the H-shaped prothoracic plate. Last instar larva came out of the seed 34.1 ± 1.59 days after oviposition (Table 1). It wandered among the seeds and pods in search of suitable place for cocoon formation. This stage lasted for 4–6 h only. Some last instar larvae were attacked by a parasitoid, *Eurytoma raoi* Narendran (Hymenoptera: Eurytomidae) and killed during development inside the cocoon. Cocoon was formed either outside a pod or inside the pod among the seeds.

Occlusion

Adult insect emerged out after cutting an incomplete circular lid at one end of cocoon and a complete circular window in the pods of the host plant. The pupae in the cocoons formed in October did not produce adult insects immediately but overwintered as pupal instar till next April. Therefore, pupal duration inside the cocoon lasted for 55.8 ± 3.01 days during favourable period (April to September) and 205.4 ± 8.82 days during unfavourable period (October to March). Life cycle of *C. crineus* was completed in 89.9 ± 3.44 days and total longevity of the insect (including egg, larval, pupal and adult stages) lasted 109.3 ± 3.71 days under natural atmospheric conditions (24°C – 39°C temperature and 54%–98% relative humidity) during the favourable period.

DISCUSSION

In *Caryedon crineus* mating and egg laying occurred during night time similar to *C. serratus* (Pajni and Mann, 1979). Behaviour of gravid female to deposit single egg on healthy and unwrinkled seed and pod of host plant is comparable with *Callosobruchus maculatus* (Huignard *et al.*, 1985; Wasserman, 1985); *C. theobromae* (Thakur and Banyal, 2004); and some other species of genus *Caryedon* (Southgate, 1979).

In the absence of seeds and pods of host plant the females of *C. crineus* delayed oviposition for 3–4 days and finally laid eggs on the walls of the test tubes and petridishes but such eggs failed to hatch out similar to *Zabrotes subfasciatus* (Pajni and Jabbal, 1986). Eggs laid by *C. crineus* towards the end of oviposition period were without ooplasm like the eggs of *Callosobruchus chinensis* where size and number of eggs decreased with maternal age (Yanagi and Miyatake, 2002). Mating duration of 10–20 min and 21.8 ± 3.2 eggs laid per female in *C. crineus* are comparable with 25–40 min long mating and 26 eggs per female in *C. serratus* (Pajni and Mann, 1979).

First instar larvae of *C. crineus* hatched out from the ventral surface of the eggs and penetrated into host seed or pod with the help of a H-shaped prothoracic plate

TABLE 1. Duration and dimensions of different stages observed during the life cycle of *Caryedon crineus* (mean \pm standard deviation).

Incubation period (days)	Egg dimensions (mm)		Dimensions of first instar larvae (mm)		Egg to pre- pupal duration (days)	Pupal duration during favourable period (days)	Pupal duration during unfavourable period (days)
	Length	Breadth	Length	Breadth			
8.9 \pm 0.37	0.840 \pm 0.06	0.496 \pm 0.03	0.469 \pm 0.02	0.237 \pm 0.01	34.1 \pm 1.59	55.8 \pm 3.01	205.4 \pm 8.82

like in *C. serratus* (Pajni and Mann, 1979), *C. theobromae* (Thakur and Banyal, 2004) and *Bruchus pisorum* (Pajni and Sood, 1984). Exhibiting similarity to the larvae of *Bruchidius atrolineatus* and *C. maculatus* (Germain *et al.*, 1987) larvae of *C. crineus* do not shift from one seed to another even in the insufficiency of food contents in previously infested seed. Last instar larva of the insect under study formed cocoon outside the infested seed among the debris like *C. serratus* (Conway, 1983).

C. crineus completed its life cycle in 89.9 ± 3.4 days on *Cassia occidentalis* seeds whereas life cycle of *C. serratus* was completed in 51 days on tamarind and in 91.63 days on *Bauhinia rufescens* seeds (Halle *et al.*, 2002). Adults of *C. crineus* appeared in the field in the month of April after passing the winter as hibernated pupa similar to *B. pisorum* (Pajni and Sood, 1976). Pupal duration of *C. crineus* inside cocoon lasted for 205.4 ± 8.82 days during unfavourable period. A similar prolonged life cycle of 6–8 months during cooler months has been observed in *C. chinensis* (Yoshida *et al.*, 1986).

REFERENCES

- Arora, G. L. (1977) Taxonomy of the Bruchidae (Coleoptera) (adults) of Northwest India. *Oriental Insects Supplement* 7: 1–132.
- Arora, G. L. (1978) Taxonomy of the Bruchidae (Coleoptera) (larvae) of Northwest India. *Oriental Insects Supplement* 8: 1–48.
- Conway, J. A. (1983) Notes on the biology and ecology of the groundnut seed beetle, *Caryedon serratus* (Ol.) (Coleoptera: Bruchidae) under field conditions in Senegambia. *Tropical Stored Products Information* 45: 11–13.
- Germain, J. F., Monge, J. P. and Huignard, J. (1987) Development of two bruchids population (*Bruchidius atrolineatus* Pic. and *Callosobruchus maculatus* Fab.) infesting stored cow-pea (*Vigna unguiculata* Walp.) pods in Niger. *Journal Stored Product Research* 23: 157–162.
- Halle, D. N., Awaknavar, J. S. and Somashekhar, (2002) Biology of tamarind beetle, *Caryedon serratus* (Olivier) on groundnut and other hosts. *Insect Environment* 8: 67–69.
- Huignard, J., Leroi, B., Alzouma, I. and Germain, J. F. (1985) Oviposition and development of *Bruchidius atrolineatus* (Pic.) and *Callosobruchus maculatus* (F.) (Bruchidae: Coleoptera) in *Vigna unguiculata* (Walp.) cultures in Niger. *Insect Science and Application* 6: 691–699.
- Pajni, H. R. and Mann, B. K. (1979) Some aspects of the biology of *Caryedon serratus* (Coleoptera: Bruchidae). *Bulletin Grain Technology* 17: 43–47.
- Pajni, H. R. and Sood, S. (1976) Some observations on the biology of pea-weevil, *Bruchus pisorum* (L.) (Coleoptera: Bruchidae). *Bulletin Grain Technology* 14: 201–205.
- Pajni, H. R. and Jabbal, A. (1986) Some observations on the biology of *Zabrotes subfasciatus* (Boh.) (Bruchidae: Coleoptera). *Research Bulletin (Science) Panjab University* 37: 11–16.
- Pajni, H. R. and Sood, S. (1984) Some observations on the life cycle of *Bruchus pisorum* (L.) (Bruchidae: Coleoptera). *Research Bulletin (Science) Panjab University* 35: 185–187.
- Pierre, D. and Huignard, J. (1990) The biological cycle of *Caryedon serratus* (Boh.) (Coleoptera: Bruchidae) on one of its host plants, *Bauhinia rufescens* (Lam.) (Casalpinioideae) in a Sahelian zone. *Acta Oecologica* 11: 93–101.
- Southgate, B. J. (1979) Biology of the Bruchidae. *Annual Review of Entomology* 24: 449–473.
- Thakur, D. R. and Banyal, H. S. (2003) A study on the life history of *Caryedon crineus*. In: *National Symposium on Frontier Area of Entomological Research*, IARI: New Delhi, 127.
- Thakur, D. R. and Banyal, H. S. (2004) Life history of *Callosobruchus theobromae* (L.) (Bruchidae: Coleoptera) on *Dolichos lab lab* in the field. *Polish Journal of Entomology* 73: 161–170.

- Thakur, D. R. and Banyal, H. S. (2005) Biocontrol potential of *Eurytoma raoi*, A larval-pupal parasitoid of *Caryedon crineus* (Bruchidae: Coleoptera). In: *National Conference on Applied Entomology; Current Status, Challenges and Opportunities at MPUAT*, Udaipur, Rajasthan, India, 170–172.
- Wasserman, S. S. (1985) Oviposition behaviour and its disruption in the southern cowpea weevil, *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Journal Economic Entomology* **78**: 89–92.
- Yanagi, S. and Miyatake, T. (2002) Effects of maternal age on reproductive traits and fitness components of the offspring in the bruchid beetle, *Callosobruchus chinensis* (Coleoptera: Bruchidae). *Physiological Entomology* **27**: 261–266.
- Yoshida, T., Igarashi, H. and Shinoda, K. (1986) Life history of *Callosobruchus chinensis* (L.) (Bruchidae : Coleoptera). In: *4th International Working Conference on Stored-Product Protection*, Donahayaaae, E. and Navarro, (Eds). Tel Aviv, Israel, 471–477.

(Received 25 October 2006; accepted 23 May 2007)



Ecdysone-neurosecretion interaction during vitellogenesis in *Poeciloceris Pictus* Fabr. (Orthoptera: Acrididae)

Versha Sharma

*Department of Zoology, Dr. H.S. Gour University, Sagar (M.P.) 470003, India
Email: varsha_zoology@rediffmail.com*

ABSTRACT: Different concentrations of β -ecdysterone (No-E-2003) in solution in the newly emerged female insect from day 1 to 30 days during the genotrophic cycle stimulates vitellogenin synthesis in the adipose tissue. This was found to be the effect of 'A' type of neurosecretory cells and inhibition of its hormone secretion regressed the activity of *Corpus allatum* (CA). On the contrary, the 'B' type cells of the *pars- intercerebralis* of EDNH system remain unaffected and their secretions reach the hemolymph via *Corpus cardiacum* (cc) and acts in conjunction with ecdysone to regulate the uptake of vitellogenin in the ovaries. This uptake is rather slower in the presence of juvenile hormone (JH). It is concluded that 'B' type neurosecretory cell hormone is comparable to a parsian -Lom OMP described by Girardie *et al.* (1996a,b) in *Locusta migratoria* that influence egg development.

© 2007 Association for Advancement of Entomology

INTRODUCTION

A lot of research has been done on the role of neurosecretion and juvenile hormone (JH) on insect ovarian development (Johannsson, 1955; Nayar, 1955; Wigglesworth, 1936, 1985; Highnam, 1962a; Dogra and Tandon, 1964; Saini, 1966, 1971; Dogra and Ewen, 1970; Gilbert, 1985; Girardie *et al.*, 1996a,b). JH plays a central role in egg production (Wigglesworth, 1985; Strong, 1965a) and acts on the fat bodies to initiate vitellogenin synthesis. The regulation of ovarian development via JH has been fully elucidated by many authors (Engelmann, 1983; Gilbert, 1985) but little work has been done on hormonal homeostasis in female reproduction.

Horn (1971); Lagueux *et al.* (1977) have suggested that the ecdysteroids in the adult female might play a role in reproduction, while according to Stay and Tobe (1981) the secretion of ecdysteroids from the late stage ovaries inhibit JH synthesis.

It is also well established that fat bodies synthesized vitellogenin and the synthesis is activated in the presence of 20-hydroxyecdysone (Bohm *et al.*, 1978). Injection of micro-quantities of 20 hydroxyecdysone stimulated vitellogenin synthesis in the mosquito *Aedes aegypti* (Fallon *et al.*, 1974). A factor like egg development

neurosecretory hormone (EDNH) was found to stimulate the synthesis of both ecdysone and vitellogenin (Lea, 1972, 1982). According to (Rossignol *et al.*, 1981) oversecretion of ecdysone affected JH secretion, hence the ecdysteroids may be inhibiting CA (Fallon *et al.*, 1974; Stay and Tobe, 1981). The role of ecdysone in adult insects is rather obscure, especially when its level is low. After noticing these ambiguities, in the present studies an attempt has been made to elucidate the effect of ecdysone on the neurosecretory system including CA along with its role on vitellogenin synthesis from adipose tissue and ovarian development in *P. pictus*.

MATERIALS AND METHODS

For the present study *Poeciloceris pictus* Fabr. (Orthoptera, Acrididae) has been used. The insects were collected from *Calotropis* plant during May/June to October and fresh *Calotropis* leaves were provided daily *ad libitum*. The sixth instar nymphs took about 6 to 8 days to become adult, if fed properly.

The newly emerged adult insects were sorted out and they were injected with different concentrations of β -ecdysone from day one to thirty when they start egg laying. Various concentrations of ecdysone was injected after every 24 hrs. The tissues studied are neuroendocrine complex, ovary and adipose tissue.

The concentration of hormone used were 1 μ g, 7.5 μ g, and 15 μ g per ml of acetone. 2 μ l of each concentration (containing 2 μ g, 15 μ g, 30 μ g ecdysone each) was injected to each experimental insect of different age group from day one to 3 days, 8 days, 15 days, 20 days, 25 days, and 30 days, by Hamilton Micro-syringe. The same concentration of acetone without ecdysone was injected in to each control insect.

The neuroendocrine complex including the neurosecretory cell groups, *Corpus cardiacum* and *Corpus allatum* were examined. The adipose tissue and ovaries were also examined during growth, reproductive period and during maturation period and progress of vitellogenesis relating to endocardial activity was also co-related.

The neurosecretory cells of the pars-intercerebralis (both of A and B type NS cells) were examined with different staining methods like, Paraldehyde-fuchsin (Gomori, 1950; Halmi, 1952; Gabe, 1953) referred as PF, and Chromo Haematoxylin – Phloxine (Gomori, 1950) referred as (CHP). For making observations on A-cells, technique of Ewen (1962) and Delphin (1965) were followed. Alcian blue after permanganate oxidation, counter stained with Phloxine followed by phosphotungstic acid was followed to stain different types of neurosecretory cells. Performic acid–alcian blue technique for cystine comparing NS also was used (Adams and Sloper, 1956).

Hematoxylin–eosin staining was performed to examine the adipose tissue and ovaries. For nucleic acids, the methyl green pyronine method Bonhag (1955) (Baker and Barbara, 1955; Pearse Everson, 1960) was applied. For staining proteins methods of and Pearse Everson (1960) were used.

RESULTS

Neurosecretory Cells: Four different types of neurosecretory cells A, B, C and D types in the median group and about 8–10 lateral cells located in the lateral area

of brain could be observed. Cell types could be identified by the different staining techniques used. A and B cells showed marked secretory activity and the secretion could be seen even in their axons while in the C and D type cells no secretory activity could be noticed; but vacuoles were noticed in these cells. Further there was no neurosecretory material noticed in the axons, and hence their activity could not be correlated with reproduction.

About 80–100 different NS cells could be localized in the median group. The A-type cell could be identified by PF technique and there were about 80 cells showing positive reaction to NS. Purple inclusions were noticed by PF and Alcian blue azan technique as seen by (Ewen, 1962; Delphin, 1965). These cells showed cyclic activity and they can play a major role in moulting and reproduction. The neurosecretion reached corpus cardiacum, and some neurosecretory axons reached the corpus allatum also. B-type cells showed faintly reddish neurosecretory material and their axons terminate inside the *Corpus cardiacum* (CC). The secretory activity of A-and B-type cells showed a correlated activity with vitellogenin synthesis, ovarian development, maturation and vitellogenesis, and the correlated NS activity has been depleted in (Fig. Plates 1 and 2 Fig. 1–5).

Different concentrations of ecdysone showed marked secretory changes in all types of cells and also during vitellogenin synthesis, ovarian development, yolk synthesis and vitellogenesis.

The ovarian development in normal female

In newly emerged female, vitellogenesis progresses and egg laying takes place within 28–30 days. Vitellogenesis starts from 15 days after emergence that is being completed with in about 25 days.

In three-day-old female, the median cells of A-type have some scattered granules that indicate secretory activity. There is a lot of secretory material in the axons but only little material is noticed inside the cell. The cells of corpus allatum are closely packed but neurosecretory material is noticed in the axons reaching the *Corpus allatum*.

Within 5–8 days, most of the neurosecretory material is passed to the CC. Cells of the *Corpus allatum* are still closely packed but start increasing in size. Within 13–15 days of emergence, the neurosecretory material was found to move from the body of the cell to the CC. But by this time yolk synthesis has started and at the same time the CA is found enlarged.

Within 19–20 days of emergence blue black granules are noticed in the body of A-type cells and the *Corpus allatum* and found enlarged. In the ovary, the terminal Oocyte has grown to its full size.

After 25 days from emergence, neurosecretory (NS) material is found to be filled with in the cell but no material is noticed in the axons. Vitellogenesis was also completed during this time.

Within 29–30 days from emergence, A-type cells refilled with purple coloured material, but the axons are not prominent, size of the *Corpus allatum* found to be reduced. Within 25 to 30 days chorion formation is noticed and egg laying started.




PLATE-1

FIGURE 1. A. Section of brain of adult *P. pictus* after 15 days treated with 0.03% Ecdysone ($\times 200$) P.F. staining showing A, B and C type of neurosecretory cells with NSM in it Ax-Axons. NSM-Neurosecretory material. B. Section of corpus cardiacum of adult *P. pictus* after 15 days treated with 0.03% Ecdysone ($\times 200$) P.F. with NSM in CHP-Chromophil cells. C. Section of corpus allatum of adult *P. pictus* after 15 days treated with 0.03% Ecdysone ($\times 200$) P.F. showing material stored in the posterior portion. SH-Sheath, V = Vacuoles, NSM = Neurosecretory material, CHP = Cells of corpus allatum. D. Section of adipose tissue of adult *P. pictus* after 15 days treated with 0.03% Ecdyson showing distinct nucleus and Vacuolated Cytoplasm ($\times 100$) M.G/P. E. Section of ovary of adult *P. pictus* after 15 days treated with 0.03% Ecdyson showing vacuoles in the ooplasm ($\times 200$) M.G/P.




PLATE-2

FIGURE 2. A. Section of brain of adult *P. pictus* after 15 days treatment with 0.15% Ecdyson ($\times 200$) P.F. showing A, B and C type of cells with NSM. B. Section of corpus cardiacum of adult *P. pictus* after 15 days treatment with 0.15% Ecdyson showing NSM in it ($\times 200$) P.F. C. Section of corpus allatum of adult *P. pictus* after 15 days treatment with 0.15% Ecdyson ($\times 200$) with cells of corpus allatum and NSM = Neurosecretory material in corpus allatum in cells. D. Section of adipose tissue of adult *P. pictus* after 15 days treatment with 0.15% Ecdysone ($\times 200$) M.G./P. E. Section of ovary of adult *P. pictus* after 15 days treatment with 0.15% Ecdyson ($\times 200$) M.G./P. FE = Follicular epithelia.

In control insects, major events are the same as that in the normal insects. But in the experimental insects which were injected with different concentrations of ecdysone, the synthesis of neurosecretory material in cells was rather slow and they were filled with materials as the transport also was slower and thus neurosecretory material reaching the *Corpus allatum* was inhibited and the corpus allatum was found regressed. There was no juvenile hormone secretion. Further, there was vitellogenin synthesis as usual and uptake of vitellogenin took place. But it was rather slower than that takes place in the presence of Juvenile hormone. On the contrary the B cells show normal secretion.

The neurosecretory activity showed that the secretion of B cells and ecdysone in conjunction play a role in the uptake of vitellogenin to the ovaries for yolk synthesis and vitellogenesis. This process is rather slower than that takes place in the presence of juvenile hormone. In comparison with normal vitellogenesis, this takes place between 25–30 days. Egg laying is also delayed and takes place after 30–35 days.

DISCUSSION

The insects were injected with ecdysone and with different concentrations it was found that the ecdysone plays different roles. It stimulated the synthesis of vitellogenin in adipose tissues showed increased RNA activity and vacuole formation. These observations are similar to that described by Fallon *et al.* (1974) and Hagedorn (1983) in mosquito. The injection of ecdysteroids to cockroaches reduced the activity of *Corpus allatum* (Friedel *et al.*, 1980; Lanzrein *et al.*, 1981, Rossignol *et al.*, 1981). Stay and Tobe (1981) observed that administration of ecdysteroids inhibit the activity of *Corpus allatum* and juvenile hormone synthesis. Present studies showed that different concentrations of ecdysone in *P. pictus* have an inhibitory effect on *Corpus allatum* which is mediated through prothoracicotropic hormone, from A-type of cells whose hormone reaches there by the axons to *Corpus allatum*. Although the activity of gonadotropic hormone (JH) was inhibited, vitellogenin synthesis in adipose tissue was found stimulated even though there was slow uptake of vitellogenin and yolk synthesis noticed in the ovaries in conjunction with some other hormone, the egg development hormone, EDNH Lea (1972). Ecdysone affects the secretion and transport of A-cell hormones of the EDNH complex while the B-cells and other cells of the EDNH complex are not affected and their hormone reaches the hemolymph via the *Corpus cardiacum*. Ecdysone acts in conjunction with the hormone to help vitellogenin uptake.

Mechanism of vitellogenesis is slower in the presence of JH. Hence a delay in vitellogenesis is noticed. The maturation period and vitellogenesis take place with in 15 to 25 days in normal females while in the experimental females it takes 25 to 30 days (egg laying takes place in 35 days). The hormone of B-type NS cells which is unaffected by the ecdysone is similar to that of Lom OMP parsian described by Girardie *et al.* (1996a) in *Locusta migratoria*. In short ecdysone appears to be a raw hormonal material that acts in conjunction with other hormones showing different modifications required for the physiological requirement of the insect.

ACKNOWLEDGEMENT

The author is indebted to Professor R. S. Saini for guidance and encouragement.

REFERENCES

- Adams, C. W. M. and Sloper, J. C. (1956) The hypothalamic elaboration of posterior pituitary principles in man, the rat and dog. Histochemical evidence derived from a performic acid-alcian blue reaction for cystine. *J. Endocrinol.* **13**: 221–225.
- Adams, T. S., Thomas, J. and Kelly, Charles E. Woods (1988) Relationship between the corpus cordiacum allatum complex and varies with the Haemolymph Ecdysteroid profile in the housefly *Musca domestica*. *J. Insect. Physiol.* **34**(2): 1105–1109.
- Baker, J. R. and Barbara, M. (1955) A simple pyronine/methyl green technique. *Q. J. Microsc. Sci.* **96**: 177–179.
- Bohm, M. K., Behan, M. and Hagedorn, H. H. (1978) Termination of vitellogenin synthesis by mosquito fat body a programmed response to ecdysterone. *Physiol. Entomol.* **3**: 17–25.
- Bonhag, P. (1955) Histochemical studies of the ovarian nurse tissues and Oocytes of the milk weed bug *Oncopeltus fasciatus*. *Dallas J. Morph.* **96**: 381–440.
- Delphin, F. (1965) The histology and possible functions of neurosecretory cells in the ventral ganglia of *Schistocerca gregaria* Forskal (Orthoptera: Acrididae). *Trans. R. ent. Soc. Lond.* **117**: 167–214.
- Dogra, G. S. and Ewen, (1970) Histology of Neurosecretory system and the ventrocerebral endocrine glands of adult migratory grasshopper *Melanophus* (orthoptera). *J. Morphol.* **130**: 451–466.
- Dogra, G. S. and Tandon, (1964) Adaptation of certain histological techniques for *in situ* demonstration of the neuroendocrinal system of insects and other animals. *Anal. J. Micro. Sci.* **105**: 445–466.
- Engelmann, (1983) In: *Vitellogenesis Controlled by Juvenile Hormone in Endocrinology of Insects*, Downor, R. G H and Laufer, H. (Eds). Alan R. Liss: New York, 259–270.
- Ewen, A. B. (1962) An improved aldehyde fuchsin staining technique for neurosecretory products in insects. *Trans. Am. Microsc. Soc.* **81**: 94–96.
- Fallon, A. M., Hagedorn, H. N., Wyatt, G. R. and Laufer, H. (1974) Activation of vitellogenin synthesis in the mosquito *Aedes aegypti* by ecdysone. *J. Insect. Physiol.* **20**: 1815–1823.
- Friedel, H., Feyereisen, R., Mundall, E. C. and Tobe, S. S. (1980) The allostatic effect of 20 hydroxy-ecdysone on the adult viviparous cockroach *Diploptera punctata*. *J. Insect Physiol.* **26**: 665–670.
- Gabe, M. (1953) Sur quelques application de la coloration par la fuch sine paraldehyde. *Bull. Micr. Appliques.* **3**: 153–162.
- Gilbert, L. I. (1985) Comprehensive insect physiology. In: *Biochemistry and Pharmacology*, Kerkut, G. A. and Gilbert, L. I. (Eds). Pergamon Press: Oxford, New York.
- Girardie, J. *et al.* (1996a) Detection of vitellogenin in the Haemolymph of larval female Locusts (*Locusta migratoria*) treated with the neurohormone LOM OMP. *J. Insect Physiol.* **42**(2): 107–113.
- Girardie, J. *et al.* (1996b) LOM OMP, a putative ecdysiotropic factor for the ovary in *Locusta migratoria*. *J. Insect Physiol.* **42**(3): 215–227.
- Gomori, G. (1950) Aldehyde-Fuchsin: a new stain for elastic tissue. *Amer. J. Clin. Path.* **28**: 665–666.
- Hagedorn, H. H. (1983) The role of ecdysteroids in the adult insects. In: *Endocrinology of Insects*, Alan R. Liss. Inc.: New York, 271–304.
- Halmi, N. S. (1952) Differentiation of two types of basophils in the adenohypophysis of the rat and mouse. *Stain. Tech.* **27**: 61.

- Highnam, K. C. (1962a) Neurosecretory control of ovarian development in *Schistocerca gregaria*. *J. Micros. Sci.* **130**: 57–72.
- Horn, D. H. S. (1971) The ecdysones. In: *Naturally Occurring Insecticides*, Jacobson, M. and Grosely, D. G. (Eds). Deyker Press: New York, 33–439.
- Johannsson, A. S. (1955) The relationship between corpora allata and reproductive organs in starved female *Leucophaea maderae* (Blattario). *Bio. Bull.* **108**: 40–44.
- Kailson, P., Stamm, and Menendez, D. (1956) Notiz über den Nachweis Von metamorphose hormon in den Imagines von *Bombyx mori* Hoppe-seyler's *Z. Physiol. Chem.* **306**: 109–11.
- Kindle, H., König, R. and Lanzrein, B. (1985) An *in vitro* system to study the incorporation of vitellogenin into the oocytes of the ovoviparous cockroach. *Nauphoeta cinerea*, *Experientia* **41**: 1225.
- King, D. S. and Marks, E. P. (1974) The secretion and metabolism of α -ecdysone by the cockroach (*Leucophaea maderae*) tissue *in vitro*. *Life Sci.* **15**: 147–154.
- Lagueux, M., Hirn, M. and Hoffmann, J. A. (1977) Ecdysone during ovarian development in *Locusta migratoria*. *J. Insect Physiol.* **23**: 105–119.
- Lanzrein, B., Wilhelm, R. and Buschor, J. (1981) On the regulation of the corpora allata activity in adult females of the ovoviparous cockroach *Nauphoeta cinerea*. In: *Juvenile Hormone Biochemistry*, Pratt, G. E. and Brooks, G. T. (Eds). Elsevier: Amsterdam, 147–160.
- Lea, A. O. (1972) Regulation of egg maturation in the mosquito by the neurosecretory system, the role of the *Corpus cardiacum*. *Gen. Com. Endocrinol. Suppl.* **3**: 602–608.
- Lea, A. O. (1982) Artificial stimulation of vitellogenesis in *Aedes aegypti* by 20-hydroxyecdysone. *J. Insect Physiol.* **28**: 173–176.
- Nayar, K. K. (1955) Studies on the neurosecretory system of *Iphita limbata* stal. Distribution and structure of the neurosecretory cells of the nerve ring. *Bio. Bull.* **108**: 296–307.
- Pearse Everson, A. G. (1960) *Histochemistry, Theoretical and Applied*, G & A. Churchill: London.
- Rossignol, P. A., Feinsod, F. M. and Spielman, A. (1981) Inhibitory regulation of corpus allatum activity in mosquitoes. *J. Insect. Physiol.* **10**: 651–654.
- Saini, R. S. (1966) Neuroendocrine control of oocyte development in the beetle *Aulacophora foveicollis*. *Luc. J. Insect Physiol.* **12**: 1003–1008.
- Saini, R. S. (1971) Neuroendocrine control of oocyte development in *Poecilercera picta*. *J. Zool. Lond.* **165**: 275–83.
- Stay, B., Friedel, T., Tobe, S. S. and Mundall, E. C. (1980) Feedback control of Juvenile hormone synthesis in cockroaches possible role of ecdysterone. *Science* **207**: 898–900.
- Strong, L. (1965a) The relationship between the brain *Corpora allata* and oocyte growth in the Central American locust *Schistocerca* sp. II. The innervation of corpora allata, the lateral neuro secretory complex and oocyte growth. *J. Insect Physiol.* **11**: 271–280.
- Tobe, S. S. and Stay, B. (1980) Control of juvenile hormone biosynthesis during the reproductive cycle of a viviparous cockroach. III effect of denervation and age on compensation with unilateral allatectomy and supernumerary *Corpora allata*. *Gen. Comp. Endocrinol.* **40**: 89–98.
- Stay, B. and Tobe, S. S. (1981) Control of the *Corpora allata* during a reproductive cycle in a viviparous cockroach. *Am. Zool.* **21**: 663–674.
- Wigglesworth, V. B. (1936) The function of the *Corpus allatum* in the growth and reproduction of *Rhodnius prolixus* (Hemiptera). *Q. J. Micro. Sci.* **79**: 91–121.
- Wigglesworth, V. B. (1985) Historical perspectives comprehensive insect physiology. In: *Biochemistry and Pharmacology*, Kerkut, G. A. and Gilbert, L. I. (Eds). Pergamon press: Oxford, New York.

(Received 27 December 2006; accepted 11 April 2007)



Study of beetles (Coleoptera) associated with stored products in Kolkata, India

1. Genus *Carpophilus* Stephens (Nitidulidae)

P. K. Basak¹ and T. K. Pal²

¹Department of Zoology, Bidhannagar College, Kolkata 700064, India
Email: basak.prabir@rediffmail.com

²Zoological Survey of India, M-Block, New Alipore, Kolkata 700053, India
Email: tkpal51@rediffmail.com

ABSTRACT: Six species of *Carpophilus* Stephens are reported from 18 different commodities. Of these, *Carpophilus freemani* Dobson, *C. obsoletus* Erichson, *C. pilosellus* Motschulsky, *C. delkeskampi* Hisamatsu and *C. marginellus* Motschulsky are new records from stored products in India. Diagnostic features, distribution and habitat records of all six species are dealt with and a key to these species of *Carpophilus* is given. © 2007 Association for Advancement of Entomology

KEYWORDS: Nitidulidae, *Carpophilus* Stephens, stored product pests

INTRODUCTION

Carpophilus Stephens is a large genus, comprising about 193 species (Jelinek, personal communication, 2006) that includes most of nitidulid species infesting stored products. This genus can be recognized by the following characters: head distinctly narrower than pronotum, clypeus slightly indistinct and margined by a depression on either side; eyes large; antennae little longer than head with compact, flattened, rounded or oval club; labrum bilobed; pronotum about as broad as elytra; scutellum broadly rounded posteriorly; elytra short, exposing two apical abdominal segments. Species of *Carpophilus* Stephens infesting stored products, especially in Europe, were dealt with earlier by Dobson (1954a,b, 1956, 1960). Grouvelle (1913) mentioned four species of *Carpophilus* from India, namely, *Carpophilus dimidiatus* (Fabricius), *C. hemipterus* (Linnaeus), *C. ligenus* Murray and *C. mutilatus* Erichson encountered commonly in stores. Aitken (1975) reported these four species intercepted from Indian products imported to U.K. and altogether nine species of *Carpophilus* from the world. In the present study, five more species namely, *C. freemani* Dobson, *C. marginellus* Motschulsky, *C. obsoletus* Erichson, *C. pilosellus* Motschulsky and *C. delkeskampi*

Hisamatsu, besides *C. hemipterus* (Linnaeus) were detected during the course of an intensive survey of stored product beetles from Metropolitan Kolkata during 1982–85. Systematic accounts of these species are given and a key to their identification is incorporated.

SYSTEMATIC ACCOUNT

Family Nitidulidae

Subfamily Carpophilinae

Genus *Carpophilus* Stephens

Carpophilus Stephens, 1830, *Ill. British Insects*, **3** 50.

Carpophilus delkeskampi Hisamatsu

Carpophilus hemipterus (nec. Linnaeus) Stebbing, 1914, *Indian Forest Insects, Coleoptera* 108.

Carpophilus delkeskampi Hisamatsu, 1963, *Entomon. Rev. Japan*, **15**: 61.

Diagnosis

Brownish yellow with paler legs, elytra yellowish with patches of brown near base, sparsely and finely pubescent (Fig. 1).

Size: 2.50–2.65 mm.

Material studied

8 exs. India: W. Bengal, Kolkata, Sovabazar, *ex.* maize grain.

19.iii.1982; P. K. Basak; 2 exs, Kolkata, Posta, *ex.* cassia leaf, 2.vii.1982; P. K. Basak; 3 exs. Howrah, Bandhaghat, *ex.* cotton, 17.11.1982, 1 ex, Kolkata, Chitpur, *ex.* jute thread 23.vi.1983, P. K. Basak. 2 exs.

Distribution

India: West Bengal (new record); Sri Lanka.

Remarks

There is no previous record of habitat of this species and in the present study it has been recorded from maize grain, cassia leaf, cotton and jute thread.

***Carpophilus freemani* Dobson**

Carpophilus freemani Dobson, 1956. *Entomologist's mon. Mag.*, **92**: 41.

Diagnosis

Brownish black with paler elytra and moderately long and recumbent pubescence (Fig. 1(2)).




FIGURE 1. 1. *Carpophilus delkekampi* Hisamatsu, Dorsal view. 2. *Carpophilus freemani* Dobson, Dorsal view. 3. *Carpophilus hemipterus* (Linnaeus), Dorsal view. 4. *Carpophilus marginellus* Motschulsky, Dorsal view. 5. *Carpophilus obsoleus* Erichson, Dorsal view. 6. *Carpophilus pilosellus* Motschulsky, Dorsal view.

Size: 2–23 mm.

Material studied

9 exs. India. Howrah, Railway yard, ex. tamarind 17.ii.1982; P. K. Basak; 7 exs., Howrah, ex. fish baskets and gunny sacs, 15.i.1982; P. K. Basak; 2 exs.

Distribution

India: West Bengal; England; North America; South America.

Remarks

Dobson (1956) described this species from specimens collected from Brazil nuts. It also infested many kinds of moist decomposing vegetable matter, including apple and peach drops, corn ear refuse and weevil infested nuts (Connell, 1956; Aitken, 1975) recorded this species from Brazil nuts, maize, bones, and sheanuts. In the present study it was recorded mostly from tamarind and less frequently from baskets and gunny sacs used to carry fishes.

Carpophilus hemipterus (Linnaeus)

Dermestes hemipterus Linnaeus, 1758. *Syst. Nat.*, ed. 10: 358.

Carpophilus hemipterus: Stephens, 1830, *Illus. Brit. Ins.*, 3: 50.

Diagnosis

Dull or shiny black, with two conspicuous, amber-brown spots at posterior tips of elytra and two smaller, more obscure spots of same colour near bases of lateral margins of elytra; antennae and legs amber coloured; surface of body is finely punctate, each pit gives rise to a hair (Fig. 1(3)).

Size: 3 mm.

Material studied

30 exs. India: Kolkata, Ultadanga, ex. keora and mahua seeds, 14.v.1983; P. K. Basak; 20 exs, Kolkata, Shyambazar, ex. groundnut seeds, 19.iii.1983; P. K. Basak; 10 exs; Kolkata, Mechuabazar, ex. walnuts, 9. iv.1983; P. K. Basak; 7 exs; Kolkata, Sovabazar, ex. maize grain, 19.iii.1982; P. K. Basak; 3 exs.

Distribution

World-wide; occurring throughout tropical and temperate parts of the world (Hinton, 1945). From India this species has been recorded from West Bengal and Bihar.

Remarks

Hinton (1945) recorded this species from grapes, raw sugar, figs, cloves, copra, dried apples, apricots, prunes, avocado, raisins, bananas, limes, peaches, plums, dates, oranges, pears, melons, tomatoes, honey, sorghum, pineapple, corn, wheat, rice bran, corn meal, cotton seeds, oats, shelled peanuts, and other nuts. This species also feeds on the pulp of the fruit itself and infests readily if contaminated with fungus. Normally they cannot enter the fruit unless the surface is broken or damaged. It was also found, in less number, on cereals, oilseeds and their products. They infest mostly 'keora' seeds, mahua and groundnut seeds and less so walnuts and maize grain.

***Carpophilus marginellus* Motschulsky**

Carpophilus marginellus Motschulsky, 858, *Etud. Ent.*, 7: 40.

Diagnosis

Dark brown, dorsally and underside reddish brown, strongly punctate, covered by recumbent golden hairs (Fig. 1(4)).

Size: 2.2–3.5 mm.

Material examined

20 ex. India: Howrah, Railway yard, ex. bentonite powder, 24.iv.1983; P. K. Basak; 5 exes., Kolkata, Sealdash Railway yard, ex. portland cement, 4.viii.1983; P. K. Basak; 2 exs., Kolkata, Sealdah, ex. tamarind, 15.vii.1983; P. K. Basak; 4 exs., Kolkata, Burrabazar, ex. wheat bran, 26.vi.1982; P. K. Basak; 3 exs., Kolkata, Posta, ex. flour, 24.iv.1985; P. K. Basak 1 ex., Kolkata, Sovabazar, ex. maize grain, 19.iii.1982; P. K. Basak; 5 exs.

Distribution

Widely distributed in tropical and sub-tropical parts of the world, including USA, Britain, W. Africa, Madagascar, Sri Lanka, Indo-China, Malayasia, Hong Kong, Japan, Pacific Islands, and presently recorded from India (W. Bengal) for the first time.

Remarks

Dobson (1954a,b) recorded this species from cocoa beans in W. Africa, and wheat from Maryland, USA. Aitken (1975) recorded it from wheat, maize, cocoa beans, soya beans, desiccated cocoanut, copra and what from different countries, and Hinton (1945) encountered it in rice and flour. In the present study, it was collected mostly from bentonite powder, moderately from tamarind, wheat bran, flour, maize grain, and occasionally from portland cement. The specimens collected from mango seeds (ZSI) have also been examined.

***Carpophilus obsoletus* Erichson**

Carpophilus obsoletus Erichson, 1843. *German Zeitschr. Fur. Ent.* 259.

Diagnosis

Blackish; elytra dark brown with pale humeral angles; sparsely pubescent (Fig. 1(5)).

Size: 2.3–3.8 mm.

Material examined

9 exs. India: Kolkata, posta, *ex.* flour, 24.iv.1985; P. K. Basak; 2 exs., Kolkata, Sovabazar, *ex.* maize grain, 19.iii.1982; P. K. Basak; 3 exs., Kolkata, Ultadanga, *ex.* mahua seeds, 14.v.1983; P. K. Basak; 2 exs., Kolkata, Shyambazar, *ex.* groundnut seeds, 19.iii.1983; P. K. Basak; 1 ex.

Distribution

Widespread throughout the tropics and sub-tropics; found in Europe, Asia, Africa including Madagascar, India, Sri Lanka, Phillipines, Malaysia, Japan, Formosa, Iraq, Solomon islands, S. America, West Indies. In India, this species is known from West Bengal, Bihar, Arunachal Pradesh, Maharashtra and Andaman Islands.

Remarks

This is a widespread pest of grains, grain products, oilseeds, copra, oilcakes, cocoa beans, dried fruits, nuts, illipenuts, and sago flour (Aitken, 1975). Sengupta *et al.* (1984) recorded it from dried plums. In the present study, this species was observed mostly in flour and maize grains, and moderately in mahua, groundnut seeds, coir etc.

Carpophilus pilosellus Motschulsky

Carpophilus pilosellus Motschulsky, 1758, *Etud. Ent.* 7: 41.

Diagnosis

Reddish brown, surface shiny with recumbent pubescence (Fig. 1(6)).

Size: 1.8–2.9 mm.

Material examined

5 exs. India: Kolkata, Ultadanga, *ex.* keora seeds, 14.v.1983; P. K. Basak; 2 ex., Kolkata, Shyambazar, *ex.* groundnut seeds, 19.iii.1983; P. K. Basak; 1 ex., Kolkata, Mechuabazar, *ex.* fruit basket, 9.iv.1983; P. K. Basak; 1 ex., Kolkata, Howrah, *ex.* fish baskets and gunny sacs, 15.i.1982; P. K. Basak; 1 ex.

Distribution

China, Japan, Madagascar, Indo-China, Sulawesi, Micronesia. From India this species has been reported from Delhi, Jharkhand and West Bengal.

Remarks

This species was recorded from ripe banana, copra, chicken feed, decaying cocoanut, leaf litter, fodder like plant, spider web and chocolate bar (Gillogy, 1962). Aitken (1975) collected it from rice, rice bran, wheat mixed with maize, brazil nuts, maize meal, oilcakes, gallnuts and illipenuts. In the present study, it was collected mostly from decaying keora seeds, and occasionally from groundnuts, fish scales, fruit baskets, and gunny sacs used for carrying fish.

KEY TO SPECIES OF *CARPOPHILUS* STEPHENS FROM STORED PRODUCTS

1. Apex of pygidium bluntly rounded 2
 - Apex of pygidium points 3
2. Pronotum 1.5x wider than long, broadest in middle, sides feebly arcuate, without basal foveal; antennal segment 3 as long as segment 2 (Fig. 1(2)) *Carpophilus freemani* Dobson.
 - Pronotum slightly wider than long, broadest in basal half, sides curved uniformly in anterior half, with basal foveae; antennal segment 3 shorter than segment 2 (Fig. 1(3)) *Carpophilus hemipterus* (L.)
3. Posterior margin of pronotum distinctly curved; antennal segment 3 shorter than or as long as segment 2 4.
 - Posterior margin of pronotum feebly sinuate; antennal segment 3 slightly longer than segment 2 (Fig. 1(4)) *Carpophilus marginellus* Motschulsky.
4. Propygidium and pygidium elongate, elytra feebly transverse with no depression at base 5.
 - Propygidium and pygidium short and transverse; elytra much wider than long and depressed at base (Fig. 1(1)) *Carpophilus delkeskampii* Hisamatsu.
5. Body oblong-oval, not shining, elytra blackish-brown with pale humeral angles; antennal club with rounded apex (Fig. 1(5)) .. *Carpophilus obsoletus* Erichson.
 - Body oblong, shining, elytra uniformly reddish brown; antennal club with acuminate apex. (Fig. 1(6)) *Carpophilus pilosellus* Motschulsky.

DISCUSSION

A large number of beetles are of considerable economic importance as they infest and damage a wide range of stored food and other commodities of biotic origin. At least 500 species have hitherto been listed as primary infesters of many plant and animal products in stores and warehouses from the world. Many more species are encountered in storage and in packing material as associates which are otherwise scavengers, predators or mould feeders. All these species have specialized ecological requirements and life patterns that influence the degree of economic loss. Identification of these beetles is quite often not easy. A survey of storage products in Kolkata was therefore undertaken to study the species found in these habitats and at least 85 species have

been found. This study of *Carpophilus* species resulted in recording 5 more species from the storage sites in India.

ACKNOWLEDGEMENTS

We are thankful to the Director, Zoological Survey of India for providing laboratory facilities. We express sincere gratitude to Dr. J. Jelinek, Department of Entomology, National Museum, Czech Republic, for providing pertinent literature and required information and to Dr. D. G. H. Halstead, formerly of Slough Laboratory, U.K. for confirming identification of several specimens. One of us (PKB) is grateful to the Principal, Bidhannagar College for encouragement. We acknowledge indebtedness to the reviewer of the paper who has critically gone through it to offer useful changes for its improvement.

REFERENCES

- Aitken, A. D. (1975) *Insect Travellers, Vol. 1. Coleoptera, Tech. Bulletin 31*, MAFF Publ.: London. **16**, p. 191.
- Connell, W. A. (1956) Nitidulidae of Delaware. *Bull. Univ. Delaware* **318**: 1–67. (Technical)
- Dobson, R. M. (1954a) The species of *Carpophilus* Stephens (Col. Nitidulidae) associated with stored products. *Bull. Ent. Res.* **45**: 389–402.
- Dobson, R. M. (1954b) A new species of *Carpophilus* Stephens found on stored products. *Entomologist's Mon. Mag.* **90**: 299–300.
- Dobson, R. M. (1956) A new species of *Carpophilus* Stephens (Col. Nitidulidae) associated with stored products. *Entomologist's Mon. Mag.* **92**: 41–42.
- Dobson, R. M. (1960) Notes on the taxonomy and occurrence of *Carpophilus* Stephens (Col. Nitidulidae) associated with stored products. *Entomologist's Mon. Mag.* **95**(1959): 156–158.
- Gillogy, L. R. (1962) Insects of Micronesia, Coleoptera: Nitidulidae. *Insects of Micronesia* **16**: 133–188.
- Grouvelle, A. (1913) Nitidulidae. *Coleopt. Cat. Part* **56**: 8–223.
- Hinton, H. E. (1945) A monograph of the beetles associated with stored products. *Brit. Mus. (Nat. Hist.), London* **1**: 443.
- Sengupta, T., Mukhopadhyay, P. and Sengupta, R. (1984) Major Beetle Pests of Stored Food Products in India. *Occ. Pap. No., Rec. Zool. Surv. India* **62**: p. 65.

(Received 6 December 2006; accepted 11 April 2007)



Impact of IPM on natural enemies in irrigated cotton of North India

R. K. Tanwar, O. M. Bambawale, P. Jeyakumar, A. Dhandapani, Vikas Kanwar, O. P. Sharma and D. Monga*

National Centre for Integrated Pest Management, LBS Building, IARI Campus, New Delhi 110012, India

Email: tanwaripm@rediffmail.com

ABSTRACT: Large-scale farmers' participatory field validation trials were conducted at village Panihari in Sirsa district of Haryana for 3 crop seasons during 2002–04 to evaluate impact of Integrated Pest Management (IPM) on conservation of natural enemies in irrigated cotton. IPM module consisted mainly of balanced fertilizer use, regular scouting and pheromone monitoring, need based application of neem products alternating with chemical insecticides and release of *Beauveria bassiana* and *Trichogramma chilonis*. The Farmers' Practices (FP) relied exclusively on chemical insecticides and use of higher/lower fertilizer dose than the recommended dose. The average number of chemical insecticidal sprays was substantially reduced to 2–3 sprays in IPM against 6–7 single or mixed sprays in FP. The overall reduction of pesticide use over three seasons reflected in high population of natural enemies i.e. 0.35 eggs + larvae of *Chrysoperla carnea* and 0.16 adults of spiders per plant in IPM compared to 0.14 eggs + larvae of *C. carnea* and 0.07 adults of spiders per plant in FP. Better parasitization of *Helicoverpa armigera* eggs by *T. chilonis* was in IPM block (22.4%) as compared to natural parasitization (2.2%) in FP.

© 2007 Association for Advancement of Entomology

KEYWORDS: Integrated pest management, *Helicoverpa armigera*, *Trichogramma chilonis*, conservation, farmers' practices

INTRODUCTION

Indiscriminate use of broad spectrum insecticides to contain the insect pests in cotton ecosystem has very often resulted development of high level of resistance in bollworms, resurgence of sap sucking insects (Dhawan and Simwat, 1997) and destruction of beneficial natural enemies. Though significant progress has been made in central and south India to reduce pesticide load by popularizing IPM technology in rainfed cotton among farmers (Singh *et al.*, 2002), management of insect pests

*CICR Regional Research Station, Sirsa (Haryana)

in North India is still largely based on the use of chemical insecticides, which has resulted in the wiping out of natural enemies. Attempts have also been made under Punjab conditions to develop and evaluate bio intensive IPM module in small plots (Verma and Gill, 1992; Brar *et al.*, 2002). During 2002–2004 large-scale farmers' participatory field trials were conducted on synthesis and validation of IPM in irrigated cotton at village Panihari (District Sirsa, Haryana) by National Centre for Integrated Pest Management (NCIPM), New Delhi.

MATERIALS AND METHODS

Field trials were conducted at farmers' fields in two blocks of 2 ha each in 2002 and 2003 and 8 ha each in 2004 to study the impact of IPM on population of important insect pests and natural enemies in irrigated cotton at village Panihari (Sirsa, Haryana). One block involved the common farmers' practices (FP) prevailing in the area to contain the insect pests whereas the other block involved the IPM practices. During 2002 IPM as well as FP included *Desi* cotton (*Gossypium arboreum*) variety, RG-8 and American (*G. hirsutum*) variety, H-1117 with standard agronomic practices. During 2003 and 2004 the trial was conducted with cultivar H-1098 in both IPM and FP blocks. In FP the farmers applied higher or lower dose of fertilizer than the recommended dose i.e. 70–80 kg urea/acre. The FP consisted entirely of sprays of chemical insecticides either alone or in mixtures whereas the IPM involved alternating application of 5% Neem seed powder extract (NSPE) or neem oil (2.5 l/ha) with 2–3 sprays of chemical insecticides (Table 1). The application of chemical insecticides was based upon the insect pests prevailing in the season; therefore, the schedule of chemical insecticides did not remain same during three years. In addition IPM also included the application of *Beauveria bassiana* @ 1.25 kg/ha (4×10^8 CFUs), 1 or 2 releases of egg parasitoid *Trichogramma chilonis* @ 150 000 ad/ha in August/September and installing 5 pheromone traps/ha from July to September in the field to monitor the population of *Helicoverpa armigera*. Field releases of *T. chilonis* was carried out when 2–3 moths *H. armigera* were trapped in each pheromone trap per night for consecutive 3–4 days or the eggs of *H. armigera* were available in the fields. A gap of one week before and after the release of *T. chilonis* and application of chemical insecticide was maintained. The details of different interventions have been indicated in Table 1.

Observations on population of egg and larvae of *Chrysoperla carnea* and adults of spiders were recorded at weekly interval per plant by randomly selecting 25 plants/acre. The data were analyzed statistically using paired 't' test. To record observation on parasitisation of *H. armigera* eggs by *Trichogramma*, the eggs of *H. armigera* were collected from FP and IPM blocks in the month of September and were transferred individually in glass vials (7.5 × 2.5 cm) under optimum laboratory conditions (25 ± 2.0 °C and 60% RH) for emergence of parasitoid or host larvae.

TABLE 1. Components of IPM and Farmers' practices (FP)

Month	2002		2003		2004	
	IPM	FP	IPM	FP	IPM	FP
July	NSPE + <i>B. bassiana</i>	1. Triazophos 1.5 l/ha + Alphamethrin 200 ml/ha			Neem cake 5 Q/ha	
Aug.	1. Alphamethrin 200 ml/ha NSPE 2. Imidacloprid 125 ml/ha NSPE	2. Quinalphos 2.5 l/ha	NSPE <i>B. bassiana</i>	1. Endosulfan 2.0 l/ha + Dichlorvos 750 ml/ha	Neem oil 2.5 l/ha <i>B. bassiana</i>	1. Meothrin 200 ml/ha 2. Triazophos 400 ml/ha + Alphamethrin 200 ml/ha
		3. Ethion 1.0 l/ha + Fenvalerate 500 ml/ha	1. Quinalphos 2.5 l/ha	2. Imidacloprid 125 ml/ha	1. Profenophos 2.5 l/ha+ Meothrin 200 ml/ha	3. Profenophos 2.5 l/ha+ Meothrin 200 ml/ha
	3. Quinalphos 2.5 l/ha	4. Monocrotophos 750 ml/ha + Fenvalerate 500ml/ha 5. Indoxacarb 500 ml/ha	NSPE	3. Lambda cyhalothrin 400 ml/ha + Neemexcel 4. Acephate 400 ml/ha + Alphamethrin 200 ml/ha	Neem oil 2.5 l/ha; <i>T. chilonis</i> (1 release) 2. Indoxacarb 500 ml/ha	
Sept.	NSPE + <i>B. bassiana</i>	6. Ethion 1.0 l/ha + Fenvalerate 500 ml/ha	2. Acephate 400ml/ha 3. Alphamethrin 200 ml/ha; <i>T. chilonis</i> Two sprays of NSPE at 10 days interval	5. Triazophos 400 ml/ha + Meothrin 200 ml/ha 6. Indoxacarb 500 ml/ha 7. Acephate 400 ml/ha + Fenpropathrin 200 ml/ha	<i>T. chilonis</i> (1 release)	4. Indoxacarb 500 ml/ha+ Acephate 125 gm/ha 5. Indox- acarb 500 ml/ha + Meothrin 200 ml/ha 6. Indoxacarb 500 ml/ha

NSPE-Neem Seed Powder Extract (12.5-25.0 Kg/ha as per stage of the crop);

Bb- *Beauveria bassiana* @ 1.25kg/ha (4×10^8 CFUs); *T. chilonis* @ 150 000 ad/ha




FIGURE 1. Population of *Chrysoperla carnea* (eggs) in IPM and Farmers' Practices (2002–04).

RESULTS AND DISCUSSION

As many as 274 species of biological control agents mainly arthropod parasitoids (162) and predators (90) have been reported (Lingappa *et al.*, 2001) but only a few bio-agents are available throughout the season in North India. In the present investigation the study was focused on the population dynamics of *C. carnea*, *Trichogramma* spp. and spiders during three consecutive years. As per the results for three years (Table 1), the average eggs and larval population of *C. carnea* remained 0.35/plant in IPM against 0.14/plant in FP. The differences were statistically significant in 2003 and 04. Weekly observations indicated highest population of the predator on 30th and 31st (0.7 eggs/plant), 38th (1.06 eggs/plant) and 31st and 32nd (0.54 egg/plant) Standard Meteorological Weeks (SMW), during 2002, 2003 and 2004, respectively (Fig. 1). During 2002 and 2004, after the initial high population build-up in 31st/32nd SMW, there was a decline in the population of eggs and larvae of *C. carnea* both in IPM and FP blocks. In FP 6–7 sprays of insecticide mixture were done (including 10–12 chemical insecticides), compared to 2–3 sprays in IPM, therefore, the reduction in the population in FP was very high and reached to a level of zero in 35th and 36th SMW in 2002 (Fig. 1). Among the three years the highest population (> 1 egg or larvae/plant) was observed in 38th Standard Meteorological week (SMW) in September, 2003. Studies made at Guntur under the field conditions on the effect of common as well as new insecticides of different origin against whitefly in cotton fields indicated significant reduction in the populations of coccinellids and *Chrysopa* compared to control (Venugopal Rao *et al.*, 1990). High level of toxicity of carbaryl (Hassan *et al.*, 1987), monocrotophos and quinalphos (Singh and Verma, 1986; Srinivasan and




FIGURE 2. Population of spiders (adults) in IPM and Farmers' Practices (2002-04).

TABLE 2. Population (number/plant) of *Chrysoperla carnea* and spiders in IPM and Farmers' fields (FP)

Block	<i>Chrysoperla carnea</i> (eggs+larvae)				Spiders (Adults)			
	2002	2003	2004	Mean	2002	2003	2004	Mean
IPM	0.31	0.38	0.35	0.35	0.19	0.08	0.11	0.13
FP	0.08	0.19*	0.16*	0.14	0.11	0.06	0.05*	0.73

*Paired 't' test significant at $p < 5\%$

Sundara Babu, 2000) and most of organophosphate insecticides (Toda and Kashio, 1997) to *C. carnea* have also been reported. Nymphs of *C. carnea* have a remarkable tolerance of pyrethroids, attributable to detoxification by pyrethroid esterase (s) (Ishaaya and Casida, 1981).

In case of adult spiders variation in population was recorded in each year and the population, in general, remained higher in IPM as compared to FP (Table 2). Low population of spiders in FP could be due to more application of chemical insecticides which are having negative effect on the population densities and species diversity of field spiders (Paik and Hwang, 1990; Lee *et al.*, 1993). Results of three years indicated highest population of spiders in IPM during 2002. In this year there were two peaks i.e. 30th and 41st SMW which showed higher population of adult spiders i.e. 0.40 and 0.46 adult/plant, respectively (Fig. 2). Among different spiders the prominent spiders observed were lynx spider, *Oxyopes javanus* (Thorell); orb spider, *Argiope minuta* (Karsh); wolf spider, *Lycosa pseudoannulata* (Boesenberg and Strand); long-jawed

spider, *Tetragnatha javana* (Thorell); *Neoscona theisi* (Walckenaer) and *Peucetia viridana* (Stoliczka).

The results on parasitisation of *H. armigera* eggs by *Trichogramma* sp. indicated 21.2 and 23.7 per cent parasitisation in IPM block against 2.1 and 1.8 per cent in FP in 2003 and 04, respectively (Table 3). The increase in the egg parasitisation by *Trichogramma* sp. could be due to field release of *T. chilonis* (one release in 2003 and two releases in 2004) and less number of chemical insecticides in IPM as compared to FP. The neem products applied in the present study have also been reported to be safe to parasitoids and predators and non-insect predators (Singh and Singh, 1996). The percentage of unhatched eggs of *H. armigera* (Table 3) also remained high in FP than IPM. This could be due to ovicidal action of many pesticides used in FP. Under the Punjab conditions Dhawan and Simwat (1997) observed that the parasitisation of *Helicoverpa* eggs declined from 4.6 in 1975 to only 0.78 per cent in 1995. In the same state recent studies conducted on evaluation of bio-intensive module (Brar *et al.*, 2002) indicated 10.7 % parasitisation of *Helicoverpa* eggs in bio-intensive module against 0.7% in insecticidal treatment.

It is evident from the above results that field application of chemical insecticides can be reduced to a greater extent by use of *neem* based plant products and field releases of bioagents. IPM tactics have definitely helped in conservation of natural enemies in irrigated cotton ecosystem of North India where the natural enemies have been severely affected by repeated large scale of exclusive use of chemical insecticides.

REFERENCES

- Brar, K. S., Sekhon, B. S., Singh, J., Shenhmar, M. and Singh, J. (2002) Biocontrol based management of cotton bollworms in Punjab. *J. Biol. Control* **16**(2): 121–124.
- Dhawan, A. K. and Simwat, G. S. (1997) Insecticide induced resurgence of sucking pests in upland cotton in Punjab, India. In: *Third International Congress of Entomological Sciences*, National Agricultural Research Center: Islamabad.
- Hassan, S. A., Albert, F., Bigler, F., Blaisinger, P., Bogenschutz, H. and Boller, J. J. (1987) Results of the third joint pesticide-testing programme by IOBC/WPRS working group Pesticides and beneficial organisms. *Journal of Applied Entomology* **103**: 92–107.
- Ishaaya, I and Casida, J. E. (1981) Pyrethroid esterase(s) may contribute to natural pyrethroid tolerance of larvae of the common green lacewing. *Environmental Entomology* **10**: 681–84.
- Lee, H. P., Kim, J. R. and Jun, J. P. (1993) Influences of the insecticidal application on the natural enemies and spider community. *Jour Indus. Tech. Grad. Sch. Dongguk Univ.* **1**: 295–307.
- Lingappa, S., Yadav, D. N. and Brar, K. S. (2001) Augmentative biocontrol within cotton IPM-Indian scenario. In: *Proceedings of the ICAR-CABU Workshop on Augmentative Biocontrol (June 29th to July 1st, 2000)*, Singh, S. P., Murphy, S. T. and Ballal, C. R. (Eds). CABI Bioscience, UK and Project Directorate of Biological Control (ICAR): India, 57–92.
- Paik, J. C. and Hwang, J. Y. (1990) Preliminary survey of Natural Enemies of Rice Pests for the Development of Integrated Control Programs. *Res. Report RDA* **33**: 153–171.
- Singh, A., Sharma, O. P., Lavekar, R. C., Bainbawale, O. M., Murthy, K. S. and Dhandapani, A. (2002) IPM Technology for Rainfed Cotton. *Tech. Bull.* **11**: 1–36.
- Singh, O. P. and Verma, G. C. (1986) Comparative toxicity of some insecticides to *Chrysoperla carnea* (Chrysopidae, Neuroptera) and *Trichogramma brasiliense* (Trichogrammatidae, Hy-

- menoptera), two arthropod natural enemies of cotton pests. *Agricultural Ecosystem and Environment* **15**: 23–30.
- Singh, R. P. and Singh, S. (1996) Neem in agriculture-advantages and disadvantages. In: *Recent Advances in Indian Entomology*, Lal, O. P. (Ed). APC Press: New Delhi, 67–82.
- Srinivasan, G. and Sundara Babu, P. C. (2000) Toxicity of insecticides to predatory green lacewing, *Chrysoperla carnea* Stephens (Neuroptera: Chrysopidae). *J. Biol. Control* **14**(1): 5–7.
- Toda, S. and Kashio, T. (1997) Toxic effect of pesticides on the larvae of *C. carnea*. *Proceedings of the Association for Plant Protection of Kyushu* **43**: 101–105.
- Venugopal Rao, N., Reddy, A. S. and Reddy, D. D. R. (1990) Effect of insecticides on the parasitoids and predators of cotton whitefly, *Bemisa tabaci* Genn. *J. Biol. Control* **4**(1): 4–7.
- Verma, G. C. and Gill, J. S. (1992) Manipulation of natural enemies in management of insect pests of sugarcane and cotton. *Proc. National symposium on Recent Advances in integrated pest Management*, PAU: Ludhiana, 161–162.

(Received 7 April 2007; accepted 22 June 2007)



Super parasitism in Indian lac insect, *Kerria lacca* (Kerr) and its implication on fecundity and resin producing efficiency of its two strains

K. K. Sharma*¹, Kavita Kumari¹ and Suman Lakhanpaul²

¹Lac Production Division, Indian Lac Research Institute, Namkum,
Ranchi 834010, India
Email: kewalkks@clri.ernet.in

²Botany Department, Delhi University, Delhi 110007, India

ABSTRACT: Parasitoids of lac insect inflict severe damage to the crop affecting adversely the resin yield and the fecundity of the insects, particularly during rainy season. There was 18.40% in *kusmi* and 26.00% parasitisation in *rangeeni* strain (female insects); *Aprostocetus purpureus* alone caused 7.8 and 11.8% parasitisation respectively. Up to nine parasitoids in *rangeeni* and six in *kusmi* were found from a single cell. Average reduction in resin produced by a single female due to parasitism varied between 17.25–39.80% in *rangeeni* and 25.24–37.91% in *kusmi* strain whereas, reduction in fecundity of lac insects ranged between 22.44–96.82% and 25.29–90.39% respectively for the two strains. As the number of parasitoids in each cell increased, there was a corresponding decrease in resin production and fecundity; the latter being affected more severely. © 2007 Association for Advancement of Entomology

KEYWORDS: Indian lac insect, *Kerria lacca*, resin-yield, fecundity, super parasitism, *Aprostocetus purpureus*

INTRODUCTION

Kusmi and *rangeeni* are two strains of Indian lac insect *Kerria lacca* (Kerr) which are commonly exploited for lac cultivation. Of the many problems confronting lac culture, the most serious, undoubtedly is the problem of management of pests and diseases of lac insect. Insect enemies of Indian lac insect, *Kerria lacca* (Kerr) cause considerable damage to the lac crop. The earliest reference on parasitoids of lac insects is of Stebbing (1910) who figured three, while Varshney (1976) has reported thirty different insects as parasitoids of lac insects. Two more parasitoids have been reported after that (Rizvi, 1986). Though lac predators are known to inflict severe damage to lac crop, losses due to parasitisation are also not insignificant. However, extent and

*Corresponding author

nature of damage caused by parasites is not well documented. Whatever little has been done, pertains mostly to extent of parasitisation and relative/seasonal abundance of parasitoids associated with lac insect (Glover, 1937; Srivastava *et al.*, 1984; Srivastava and Mehra, 1984). Also, Majumdar *et al.* (1962) showed interrelationship amongst population of parasitoids of lac insect, and Chowdhury *et al.* (1971) worked out relationship between density of lac insect and its parasitoids.

Fecundity of lac insect and the resin secreted by it are two important economic parameters; the former being important for propagating next crop while the later is product of commerce. Sharma and Ramani (2001) observed parasitoids reduce yield and fecundity. Instances of super parasitism are not uncommon in lac insects. *Aprostocetus purpureus* Cam. (Eulophidae) is one of the most abundant parasitoids associated with lac insects. Hence, in the present study effect of super parasitism caused by *A. purpureus* on fecundity and resin producing efficiency of Indian lac insect, *Kerria lacca* (Kerr) has been evaluated.

MATERIALS AND METHODS

Mature individual female lac insect cells at crop maturity were collected from the Institute Research Farm from lac crops on *Butea monosperma* (*palas*) for *rangeeni* and *Schleichera oleosa* (*kusum*) for *kusmi* strain during rainy season generation as lac culture is more vulnerable to parasitoids in this season. 500 cells from each host-plant were collected. Recommended package of practices for lac cultivation was followed. The cells were put individually in five ml glass vials and then plugged with cotton. After a month of storage, emergence of lac larvae and parasitoids was completed. Then the cells were grouped into healthy and parasitized ones based on the presence of small neatly cut hole(s) in the cells by emerging parasitoid(s) and/or presence of the parasitoid(s) in the cell/vial. Cells parasitized only by *Aprostocetus purpureus* were considered for grouping in to separate classes based on the number of parasitoids emerged for the present study.

Size, weight of lac secreted and fecundity of cells belonging to each class were measured after a month of storage. Since female cells are globular in shape, average of horizontal and vertical diameter measured with the help of a screw gauge was taken as cell size. Total number of larvae that emerged from the female lac insects in the vial and the larvae which could not emerge were counted after breaking the cell open. The whole count was used as fecundity of the female. Dead female insect was removed and weight of resin secreted by individual cell was recorded using an electronic balance. Healthy cells were also grouped into corresponding different classes on the basis of their size. Cells not conforming to any of the class in parasitized population were not considered for the study. Mean value of each class was used to compare various parameters of healthy and parasitized cells by paired 't' test (Panse and Sukhatme, 1985).

RESULTS

Extent of parasitisation

Rangeeni strain of lac insect was more vulnerable to parasitoids (26.0%) than *kusmi* (18.4%). Among all the recorded parasitoids of lac insect, *Aprostocetus purpureus* alone caused more than 40% parasitization in both the strains (7.8 and 10.8% respectively in *kusmi* and *rangeeni* strains). Sharma *et al.* (1997) have also recorded similar frequency of relative parasitism in lac insect. Instances of super parasitism were recorded from both the strains. Up to nine parasitoids were recorded from a single cell in *rangeeni* and six in *kusmi* strain. Frequency of occurrence of parasitized cells was indirectly related to the number of parasitoids in a cell.

Effect of super parasitism on resin yield and fecundity in *rangeeni* strain

Though mean diameter of the parasitized cells did not differ significantly from that of healthy cells, resin secreted by a parasitized lac insect was 23.38% less than the healthy cells. As the number of parasitoids in a cell increased, there was a proportional and significant decrease in the resin secreted as well as fecundity of the insect. The effect is more pronounced on fecundity. Mean fecundity per cell in a healthy insect was 325.74 in comparison to 197.78 in parasitized cells. Decrease in fecundity was recorded between 22.44%–96.82% depending upon the number of parasitoids in the cells (Table 1).

Effect of super parasitism on resin yield and fecundity in *kusmi* strain

Weighted mean of all the parameters scored viz., diameter of the cell, weight of the cell and fecundity were higher than the *rangeeni* strain. Difference in the mean diameter of the parasitized and healthy cells was insignificant, but mean of the resin secreted by an insect affected by the parasitoid was 14.567 mg which was 23.24% less than the resin secreted by the healthy insects (Table 2). Similarly, fecundity was affected much more adversely. Mean fecundity of healthy insects ranged between 281–427 in different size classes and that of parasitized cells 27–319. With increase in number of parasitoids in a cell, resin secreted by the insect and its fecundity also decreased.

DISCUSSION

Thickness of lac encrustation is one of the criteria for assessing the quality of broodlac. It is evident from the data that size of parasitized as well as healthy cells did not differ much but comparison of weight of the resin secreted by the two revealed that the amount of the resin produced by the parasitized cells was significantly lower. Hence, visual assessment of broodlac quality on the basis of encrustation thickness alone may prove to be deceptive unless weight is also taken into account.

Female lac insects are not only chief lac producers but they also serve to propagate next crop as broodlac (analogous to seed in other crops). Any adverse effect on fecundity of lac insect would have far reaching implications on the next crop. Mehra

TABLE 1. Effect of super-parasitism on fecundity and resin yield of the *rangeeni* strain of Indian lac insect, *Kerria lacca* (Kerr) during rainy season crop

Parasitoids recorded from a single cell	Cells parasitized by <i>Aprastocetus purpureus</i>						Healthy cells				
	Frequency	Mean Diameter (mm)	Mean Weight (mg)	% reduction of resin over healthy cells	Mean Fecundity	% reduction over healthy cells	Frequency	Class	Mean Diameter	Mean Weight	Mean Fecundity
1	21	3.42	10.60	17.25	280	22.44	39	3.41-3.50	3.43	12.81	361
2	15	3.34	9.43	24.80	206	43.09	42	3.31-3.40	3.36	12.54	362
3	10	3.27	8.16	33.37	155	55.71	57	3.21-3.30	3.24	12.32	350
4	6	3.07	7.69	35.92	112	65.85	50	3.01-3.10	3.06	11.74	311
5	3	3.15	7.50	35.90	97	68.81	72	3.11-3.20	3.16	12.00	328
6	1	2.92	6.89	37.71	56	79.78	35	2.91-3.00	2.95	11.25	299
7	1	2.77	7.06	37.24	93	72.24	11	2.71-2.80	2.74	10.82	215
8	1	2.69	6.01	39.80	7	96.82	5	2.61-2.70	2.66	9.98	220
9	1	2.86	6.70	37.99	30	86.05	21	2.81-2.90	2.85	10.99	277
Weighted Total	59	193.73	540.95	-	11669	-	332	-	1042.19	3972.97	108146
Weighted Mean	-	3.284 ^{ns}	9.169 ^{**}	23.38	197.78 ^{**}	39.28	-	-	3.153	11.967	325.74

ns - non significant, ** = $P < 0.01$

TABLE 2. Effect of super-parasitism on fecundity and resin yield of the *kusmi* strain of Indian lac insect, *Kerria lacca* (Kerr) during rainy season crop

Parasitoids recorded from a single cell	Cells parasitized by <i>Aprostocetus purpureus</i>						Healthy cells				
	Frequency	Mean Diameter (mm)	Mean Weight (mg)	% reduction of resin over healthy cells	Mean Fecundity	% reduction over healthy cells	Frequency	Class	Mean Diameter of the cells	Mean Weight of the cell (mg)	Mean Fecundity
1	17	3.96	15.7	25.24	319	25.79	55	3.91-4.00	3.93	21.0	427
2	9	3.85	15.0	27.54	300	27.18	61	3.81-3.90	3.84	20.7	412
3	8	3.70	13.8	29.59	212	45.50	94	3.71-3.80	3.74	19.6	389
4	3	3.64	11.9	33.15	132	60.83	50	3.61-3.70	3.65	17.8	337
5	1	3.47	9.5	37.91	27	90.39	32	3.41-3.50	3.46	15.3	281
6	1	3.52	10.6	36.90	81	73.18	45	3.51-3.60	3.55	16.8	302
Weighted Total	59	149.48	568.1	-	10323	-	337	-	1254.92	6395.7	124615
Weighted Mean	-	3.833 ^{ns}	14.567 ^{**}	23.24	264.69 ^{**}	28.42	-	-	3.724	18.978	369.78

ns - non significant, ** = $P < 0.01$

and Majumdar (1963) have studied emergence of larvae from parasitized female lac cells and concluded that emergence of larvae is not always hampered by parasitisation which is contrary to the observations of the present study, where it was found that decrease in fecundity of lac insect due to parasitisation ranged between 25.79–90.39 in *kusmi* and 22.44–96.82% in *rangeeni* strain. Thus, proportionately more broodlac would be required as compared to healthy broodlac for inoculating the same number of trees. If parasitized at an early stage, the lac insect is practically eaten up by the developing parasitoid rendering the lac useless for broodlac purpose. Moreover, the broodlac harbouring parasitoids if used for raising next crop would serve as source of infection to new lac culture.

Lac insects which are parasitized at an early stage die before reaching maturity. Also, Jaiswal and Saha (1995) have found positive and significant correlation between density of lac insects and number of parasitoids. Since the present study was confined only to isolated and adult female lac insects, it is highly likely that extent of parasitisation in continuous encrustation of lac insects under field conditions would be higher than recorded causing more damage.

ACKNOWLEDGEMENTS

Financial grant received from the Department of Biotechnology, Govt., of India for carrying out the study is thankfully acknowledged. Authors feel thankful to Dr. Bangali Baboo, Director, Indian Lac Research Institute for his encouragement and Mr. P. A. Ansari, T-4 for providing technical assistance.

REFERENCES

- Chowdhury, M. K., Sen, P. and Bhattacharya, A. (1971) Studies on the relationship between the density of lac insect, *Kerria lacca* (Kerr) (Homoptera: Coccidae) and chalcid parasitization. *Indian J. Ent.* **32**(2): 180–182.
- Glover, P. M. (1937) *Lac Cultivation in India*, Indian Lac Research Institute: Namkum, Ranchi, p. 137.
- Jaiswal, A. K. and Saha, S. K. (1995) Estimation of the population of parasitoids associated with lac insect, *Kerria lacca* (Kerr) on the basis of biometrical characters. *J. Ent. Res.* **19**(1): 27–32.
- Majumdar, N., Bhattacharya, A. and Chatterjee, A. C. (1962) Correlation coefficient between the major chalcid parasites attacking *Laccifer lacca* Kerr. *Sci. and Cult.* **28**(8): 393–394.
- Mehra, B. P. and Majumdar, N. (1963) Emergence of lac larvae from parasitised female lac insects *Laccifer lacca* Kerr. *Sci. And Cult.* **29**(4): 203–206.
- Panse, V. G. and Sukhatme, P. V. (1985) *Statistical Methods for Agricultural Workers*, Indian Council of Agricultural Research: New Delhi, p. 359.
- Rizvi, S. (1986) Studies on the taxonomy of the family aphelinidae with observations on some closely related families *Ph.D. Thesis*, Aligarh Muslim University, Aligarh.
- Sharma, K. K., Jaiswal, A. K., Bhattacharya, A., Mishra, Y. D. and Sushil, S. N. (1997) Emergence profile and relative abundance of parasitoids associated with Indian lac insect *Kerria lacca* (Kerr). *Indian J. Ecol.* **24**(1): 17–22.
- Sharma, K. Krishan and Ramani, R. (2001) Parasites effected reduction in fecundity and resin yield of two strains of Indian lac insect, *Kerria lacca*. *Indian J. Ent.* **63**(4): 456–459.

- Srivastava, D. C., Chauhan, N. S. and Teotia, T. P. S. (1984) Seasonal abundance of insects associated with the Indian lac insect, *Kerria lacca*. *Indian J. Ecol.* **11**(1): 37–42.
- Srivastava, D. C. and Mehra, B. P. (1984) Studies on the abundance of various insects associated with the Indian lac insect, *Kerria lacca* (Kerr). *Indian J. Ecol.* **7**(1): 96–104.
- Stebbing, E. P. (1910) A note on the lac insect (*Tachardia lacca*), its life history, propagation and collection. *Indian For. Mem.* **1**: No. 3
- Varshney, R. K. (1976) A check-list of insect parasites associated with lac. *Oriental Insects* **19**(1): 55–78.

(Received 8 August 2006; accepted 15 February 2007)



New species of *Thelcticopis* (Araneae: Sparassidae) from India

K. Sunil Jose* and P. A. Sebastian

Division of Arachnology, Department of Zoology, Sacred Heart College, Thevara, Kochi, Kerala 682013, India

ABSTRACT: A new species of spider belonging to the genus *Thelcticopis*, namely *Thelcticopis moolampilliensis* sp. nov., is described and illustrated from Kerala, India.

© 2007 Association for Advancement of Entomology

KEYWORDS: *Thelcticopis*, spider, Kerala, Sparassidae

INTRODUCTION

Thelcticopis is a small oriental genus of large huntsman spiders. This genus is characterized by the presence of 5–6 small teeth on the inner margin of chelicerae and 5–7 pairs of ventral spines on tibia of first leg. Spinnerets in this genus are supported on a membranous stalk and strengthened by a hairy chitinous ring. Pioneering studies on Indian *Thelcticopis* was done by Pocock (1901). He described four species from India namely *T. ajax* Pocock, 1901 from Ootacamund, *T. bicornuta* Pocock, 1901 from Naga Hills, *T. rufula* Pocock, 1901 from Nilgiri Hills, and *T. virescens* Pocock, 1901 from Trivandrum. Later *T. maindroni* Simon, 1906, *T. canescens* Simon, 1887, *T. serambiformis* Strand, 1907, *T. ancorum* Dyal, 1935 and *T. telonotata* Dyal, 1935, were reported from Indian region. Recently Sethi and Tikader (1990) published redescription of *T. canescens* Simon, 1887 and *T. maindroni* Simon, 1906. A new species of *Thelcticopis* is described and illustrated here. This is the only species of *Thelcticopis* recorded from Kerala other than *T. virescens* Pocock, 1901.

MATERIALS AND METHODS

Spiders were collected by methods described by Tikader (1987). Collected spiders were preserved in 70% ethyl alcohol and studied under Stereomicroscope. All measurements were taken with an eyepiece graticule. Epigyne was studied by clearing in 10% KOH. The status of the species was confirmed by referring to Pocock (1900,

*Corresponding author

1901) and Barrion and Litsinger (1995); it is also confirmed at Zoological Survey of India, Kolkata. The type specimens are now at the Arachnological Collections of Department of Zoology, Sacred Heart College, Thevara; which will in due course be deposited at the National Collections of Zoological Survey of India, Kolkata.

Abbreviations used are as follows: AME = Anterior median eyes, ALE = Anterior lateral eyes, PME = Posterior median eyes, PLE = Posterior lateral eyes, L = Length, W = Width, H = Height.

RESULTS

Thelcticopis moolampilliensis sp. nov.

Colour: Dorsum of cephalothorax uniformly reddish brown without any patches, legs reddish brown, a little paler than cephalothorax. Labium, sternum, maxillae reddish brown. A slightly pale dorsal line on the cephalothorax. Chelicerae reddish, fang and teeth deep reddish brown. Eyes reddish yellow, bordered by black margin. Abdomen whitish on the dorsum, lateral edges bear dark brown, broad, marginal patches, inner end of these patches uneven. Mid dorsal line bears five dark brown spots; middle three spots are more broader than anterior and posterior spots. Spinnerets reddish brown. Ventrums of abdomen white with a moderately sized patch on the middle at about the centre of epigyne and spinnerets. Tarsal claws and teeth deep reddish brown. Scopulae on labium and maxillae reddish yellow. Leg spines deep reddish brown, pedipalp reddish brown.

In living specimens, the cephalothorax is reddish brown and the abdomen white in color.

Female

Cephalothorax: Longer than wide, clothed with white pubescence. Anterior 1/5th abruptly narrowed; front margin convex with thin hairs. Fovea fine, long, longitudinal. **Eyes:** Eight in two rows encircled by black margins; anterior eye row slightly shorter and procurved, AME slightly larger, and closer to each other. Posterior eye row longer and procurved, eyes widely separated, PME much closer to each other than to laterals. Ocular quadrangle longer than wide, narrower in front. **Sternum:** Longer than wide, heart shaped, reddish brown, paler towards anterior margin, anterior end truncated, posterior end pointed, clothed with hairs, broadest at coxae I & II, and narrower towards coxae III & IV. **Labium:** Similar in colour to sternum, longer than wide, anterior half narrowed and bluntly rounded, scopulae present. **Maxillae:** Longer than broad, broader apically, scopulae brown. **Chelicerae:** Strong, outer surface more convex; inner margin with five teeth and outer margin with three slightly larger teeth. **Legs:** Reddish brown, long and stout, clothed with hairs and few spines. Leg formula 1423. Femur I with two dorsal spines, femur II, III, IV with three dorsal spines. Tarsal claws two, with nine teeth in each claw. Femur of palp with four dorsal spines, tibia with five spines; femur longer than tibia. Tibia I with five pairs of ventral spines.




FIGURE 1. *Thelcticopis moolampilliensis* sp. nov. (female) A. Dorsal view of cephalothorax and abdomen, B. Eyes, C. Sternum with maxillae and Labium, D. Epigynum ventral view, E. Epigynum, dorsal view.

Metatarsus and tarsus I with ventral scopulae. *Palp*: Femur with three, tibia with four, and tarsus with three spines.

Abdomen: Longer than wide, narrowed posteriorly, widest at the middle. Two pairs of sigilla present. Ventrums pale, a large dark brown spot near to spinnerets. *Spinnerets*: Supported by a hairy chitinous ring; anterior spinnerets widely separated. *Epigynum*: With two lateral sclerotized pieces; posterior half of which are slightly

TABLE 1. Leg measurements for *Thelcticopis moolampillyensis* ♀ (mm)

Leg	Femur	Patella	Tibia	Metatarsus	Tarsus	Total
I	4.8	1.5	4.0	3.1	1.3	14.7
II	4.5	1.4	3.5	3.1	1.2	13.7
III	3.8	1.2	3.0	3.0	1.1	12.1
IV	4.0	1.2	4.0	3.5	1.2	13.9

separated in the middle, anterior margin bordered by a narrow sclerotized margin. A widely depressed, open cavity present in the anterior half, partly divided by median segment in the middle.

Measurements: (Table 1) Total length: 17 L, Cephalothorax 6 L, Abdomen 10 L.

Ocular measurements: Eye diameter: AME = 0.36, ALE = 0.252, PME = 0.144, PLE = 0.18. Eye separation: AME-AME = 0.288, AME-ALE = 0.432, PME-PME = 0.648, PME-PLE = 0.828.

Male (Immature)

Cephalothorax: Body pattern and colour similar to female. Clypeus narrow, less than half the diameter of AME. *Eyes:* Both rows of eyes straight or very slightly procurved, posterior row longer than anterior row. PME smaller than other eyes. AME nearer to each other than ALE. *Legs:* Femur I & II without ventral spines. Tibia I & II with five pairs of ventral spines. Metatarsus I with two ventral spines on each side, femur II with five dorsal spines. Palp: Tarsal claws with four teeth, tarsus with two spines on each side. *Chelicerae:* Outer margin with three large teeth, inner margin with five smaller teeth.

Abdomen: More yellowish than female, variegated with median and lateral brownish patches. median patch consists of five spots, and lateral patches also have five spots with posterior patches being contiguous. Ventrums pale yellowish, without the black patch seen in female.

Specimen examined: Holotype 1♀, Moolampilly Is., Kochi, Kerala, India, 30.xii.2000, Coll. Sunil Jose, K.

Paratype: 2♀♀, 1♂ immature, Kalamasserry, Kochi, 20.ii.2001, Coll. Sunil Jose, K.

Distribution

Moolampilly Island and Kochi, Kerala, India.

Natural History:

Collected from leaves of *Mangifera indica* at night (8 PM) found moving along the leaves.

Diagnosis: Dorsal surface of abdomen white, with lateral edges bearing dark brown broad marginal patches. Ventrums with a small isolated black patch present little below the middle. Chelicerae with inner margin bearing five teeth and outer margin bearing three slightly larger teeth. Leg 1 with five pairs of ventral spines.

Etymology:

The species is named after the type locality, Moolampilly Island, near Kochi.

Remarks:

T. moolampilliensis is closely similar to *T. rufula* Pocock, 1901, but can be separated by following differences:

1. Tibia of first pair of legs with five pairs of ventral spines in *T. moolampilliensis* as against six pairs of spines in *T. rufula* Pocock, 1901.
2. In *T. rufula* Pocock, 1901, ventral side of abdomen with a median dorsal band consisting of a series of black spots, whereas in *T. moolampilliensis* only a small isolated black patch present little below the middle.
3. Legs, sternum and coxa ochre yellow in *T. rufula* Pocock, 1901, whereas these are reddish brown in *T. moolampilliensis*.
4. Epigyne in *T. moolampilliensis* with two lateral sclerotized pieces in the posterior half and a widely depressed, open cavity in the anterior half. Female is not so far described in *T. rufula* Pocock, 1901. Collection of the immature male of *T. moolampilliensis* helped separating it from *T. rufula* Pocock, 1901.

ACKNOWLEDGEMENTS

We thank Rev. Fr. A. J. Savianze CMI, Principal, Sacred Heart College Thevara, Kochi, for facilities and encouragement. We are also grateful to Dr. B. K. Biswas, Arachnida section, Zoological Survey of India, Kolkata for confirming the new identity of the species. Financial assistance from Kerala State Council for Science, Technology and Environment (KSCSTE) is gratefully acknowledged.

REFERENCES

- Barrión, A. T. and Litsinger, J. A. (1995) *Riceland Spiders of South and Southeast Asia*, CAB International: Wallingford, UK, xix +, p. 700.
- Pocock, R. I. (1900) *Fauna of British India*, Arachnida, Taylor and Francis London: p. 246.
- Pocock, R. I. (1901) Description of some new species of spiders from British India. *Journal of Bombay Natural History Society* 478–498.

- Sethi, V. D. and Tikader, B. K. (1990) Studies on some giant crab spiders of the family Heteropodidae from India. *Part II: Records of Zoological Survey India* **87**(2): 165–186.
- Tikader, B. K. (1987) *Hand Book on Indian Spiders*, Zoological Survey of India: Kolkata, p. 251.

(Received 31 May 2005; accepted 15 February 2006)



Activity pattern of juvenile hormone during metamorphosis in *Dysdercus cingulatus* Fabr. (Heteroptera: Pyrrhocoridae)

U. Gayathri Elayidam* and D. Muraleedharan

Centre for Arthropod Bioresources and Biotechnology (CABB),
Department of Zoology, University of Kerala, Kariavattom, Trivandrum 695581,
Kerala, India

ABSTRACT: JH being a morphogenetic hormone during nymphal stages, its titer at particular life stages determines which type of molt the developing insect has to undergo. Application of Juvenile hormone analogues on to 5th instar nymphs suggests that metamorphosis is a sequence of events each one of which is dependent on JH absence. © 2007 Association for Advancement of Entomology

KEYWORDS: juvenile hormone, methoprene, metamorphosis, *Dysdercus cingulatus*

Juvenile hormones are a homologous series of sesquiterpenoids that are involved in embryogenesis, molting, metamorphosis and reproduction (Gilbert *et al.*, 2000). In hemimetabolous insects metamorphosis involves wing growth, heteromorphic growth of segments of the legs and /or body segments. Further, the insect develops musculature necessary for flight and acquires the ability to respond to hormonal and or environmental cues involved in reproduction (Kumaran, 1990).

The insect *Dysdercus cingulatus* adults used for this study were reared in our laboratory under controlled conditions (temp: 28 ± 31 °C, L/D cycle 12:12 and r.h 90 ± 3) and fed on soaked cotton seeds. Eggs were laid in clusters between cotton seeds and were collected at various intervals, and maintained so that enough number of nymphs and adults of known age were available for the present study.

Haemolymph was collected by cutting the antennae and draining hemolymph into an eppendorf tube. Tyrosinase activity was inhibited by the addition of a few crystals of phenylthiourea. It was then centrifuged at 10,000 rpm for 10 min at 4 °C in a refrigerated centrifuge (Universal 16 R, Hettich, Zentrifugen Germany) to remove the hemocytes and other debris. Supernatant was mixed with double the volume of sample buffer.

Abdomen was cut open; fat body was taken out and washed in Insect Ringer, homogenized in a pre-chilled homogenizer with homogenizing medium and centrifuged

*Corresponding author




FIGURE 1. Electrophorogram showing changing protein profile in the hemolymph and fat body. Lane 1: Marker; Lane 2: 5th instar nymph hemolymph; Lane 3: 4-day-old adult hemolymph; Lane 4: Adult fat body; Lane 5: 0-day-old adult hemolymph; Lane 6: 5th instar fat body.

at 5,000 rpm for 10 min at 4 °C. Supernatant was mixed with equal volume of sample buffer.

The electrophoretic protein profile of haemolymph was determined by one dimensional sodium-dodecyl-sulphate polyacrylamide gel electrophoresis (SDS-PAGE) carried out according to Laemmli (1970) under discontinuous and dissociating buffer systems. Gel electrophoresis was done using a vertical slab gel mini-electrophoresis unit (Genei, Bangalore) attached to power supply (LKB-2297 macro drives). One mm thick resolving gel (1.5 M Tris HCl pH 8.8) was prepared and stacking gel (0.5 M Tris HCl pH 6.8) was laid over the resolving gel.

Protein samples were mixed with sample buffer (50 mM Tris-HCl, pH 6.8, 2% (w/v) SDS, 10% (v/v) Glycerol, 0.1% Bromophenol blue, 5% 2-mercapto ethanol) such that the protein concentration of each sample was approximately 100 µg per well. The samples were boiled for 1–2 min in water bath maintained at 100 °C. The samples were then centrifuged at 5000 rpm for 10 min in a centrifuge to remove the debris. 10 µl each was loaded on to the wells. Standard molecular markers were also run along with the samples. Electrophoresis was carried out initially at 60 V till the sample entered the resolving gel and at 120 V till the end of the run.

Juvenile hormone analogue, Methoprene (gift from Govindhan Bhaskaran, Texas A&M University, USA) was dissolved in acetone so as to get a concentration of 1 µg/µl solution. Methoprene was topically applied to the ventral region of 0-day, 2-day and 5-day old 5th instar nymphs with the help of Hamilton dispenser syringe. Control insects were treated with the same dosage of acetone (Chinzei *et al.*, 1994).

Fat body protein content of 6-day old 5th instar nymph was significantly lower than the hemolymph protein content of same age group (Fig. 1). Protein patterns of these tissues at the nymphal stage reveal the appearance of some specific bands




FIGURE 2. Electrophorogram showing changing protein profile in the hemolymph and fat body after Methoprene treatment. Lane 1: Marker; Lane 2: Treated fat body; Lane 3: Control fat body; Lane 4: Treated hemolymph; Lane 5: Control hemolymph.

that disappear after molting and certain bands that appear newly after adult eclosion (Fig. 1). Specific bands corresponding to MW between 21 kDa and 45 kDa present in the haemolymph of 5th instar nymph (6-day) were found to be absent in 0-day old adult. A reduction in the number of protein bands was also noticed in 0-day adults. But compared to 0-day adults, 6-day-old adult showed several more bands. Protein band of MW 114 kDa was more prominent in 6-day-old than that in 0-day-old adult. Unlike haemolymph proteins, different fat body protein bands of 6-day-old 5th instar nymphs were very feeble and weak and they did not vary significantly among themselves. However, all these protein bands are thin. The fat body of adult *Dysdercus* showed an increase in the number of bands compared to that in its 5th instar nymph. All the bands were prominent that showed equal staining intensity as well.

When methoprene ($0.5 \mu\text{g}/\mu\text{l}$) was applied on to just molted 5th instar nymphs, this emerged as supernumerary nymph (Fig. 3). Two-day-old 5th instar nymph that received $0.5 \mu\text{g}$ and $1 \mu\text{g}/\mu\text{l}$ developed into adultoids after subsequent ecdysis. When applied same dosage four days after the fifth instar nymphal molt, they prolonged the fifth instar stage for 2–3 days and more than 50% emerged into adults. The supernumerary nymphs and the adultoids retained varying degrees of nymphal characters. Wings are somewhat developed. However they do not cover the abdomen completely. Abdominal spots that are not present in normal adults were found in adultoids and supernumerary nymphs (Fig. 3).

Haemolymph and fat body were collected from methoprene-treated 5th instar nymph after 48 h and was subjected to SDS PAGE. Protein pattern of haemolymph




FIGURE 3. Effect of JHa on *D. cingulatus* nymphs. 1. Normal adult. 2. Normal 5th instar nymph. 3. Supernumerary nymph emerged after methoprene treatment. 4, 5, 6. Adultoids emerged after methoprene treatment showing varying deformities. 7, 8, 9. Methoprene treatment on 5-day-old 5th instar nymphs prolonged metamorphosis.

and fat body of methoprene-treated and control 5th instar nymph is as shown in Fig. 2. There is significant reduction in the intensity of almost all protein bands in the fat body of methoprene treated nymph when compared to that in the control nymph. But in the case of haemolymph protein profile, most of the proteins showed a drastic reduction in methoprene- treated nymph.

Application of JHa soon after emergence into the 5th instar nymph is found to be the most sensitive period to elicit its effect on retention of larval characters. When JHa was applied at the end of the sensitive period i.e. late fifth instar, adultoids are produced after ecdysis. Morphologically perfect extra-nymphal instars are formed when the effective concentration of the juvenoid has been maintained or surpassed during the

whole duration of the critical JH sensitive period. Different intermediate forms were found to develop when the effective juvenoid concentration has been reached for just a part of the critical period.

JHa was found to prolong the process of metamorphosis in *Trichoplusia ni* (Jones, 1985), *Xenopsylla skrjabini* (Ershova *et al.*, 1989), *Blatta Orientalis* (Short and Edwards, 1992b), *Rhodnius prolixus* (Chinzei *et al.*, 1994), *Bombyx mori* (Bharati and Miao Yungen, 2001) and *Apis mellifera* (Elekonich *et al.*, 2002). The insects which survived always completed metamorphosis by imago hatching, confirms the same results as seen in *D. cingulatus* in the present study.

Electrophoretic studies showed that there is a reduction in both fat body and haemolymph protein profile after methoprene treatment. Tiwari (1989) reported a reduction in haemolymph protein of *Diacrisia obliqua* after JHa treatment. Significant depression of total protein contents was noted in methoprene-treated pupae and adults of *Daphnia magna* when compared to its normal untreated insect (Olmstead and LeBlane, 2001). Methoprene treatment caused a significant reduction in blood protein levels of *Locusta migratoria* (Cotton and Anstee, 1991) and also in *Heteropeza pymaea* and *Mycophila speyeri* (White and Czajikowska, 2001). All these findings shows the commitment of epidermal cells to ecdysteroids that is dependent on JH activity pattern.

JH regulate the synthesis of specific larval and adult proteins in this tissue (Laufer and Borst, 1983). The mode of action of JHa lies in their stability to block the depression, transcription, or utilization of fresh genetic information (Wyatt and Davey, 1996). Juvenilizing activity of JHa is dependent on the stage of development at the time of its application. Juvenile hormone acid a catabolite of JH that induces competence of epidermal cells after ecdysteroid initiated commitment for metamorphosis (Ismail *et al.*, 1998, 2000). Studies with Juvenile hormone analogues on to 5th instar nymph suggest that metamorphosis is a sequence of events which are each dependent on the absence of JH.

ACKNOWLEDGEMENT

The authors gratefully acknowledge research grant received for this work from Kerala State Council for Science Technology and Environment, Govt. of Kerala [Project No. (T) 2 / SRS / 2004 /CSTE].

REFERENCES

- Bharati, D. and Yungen, Miao (2001) Effect of methoprene, MH-III and combinations of methoprene and MH III on larval Adult moth characters, Cocoon Quality and silk proteins of Silkworm, *Bombyx mori* L fed on Mulberry leaf and artificial diet. *PJS V* 130(1).
- Chinzei, Y., Azumi, C., Miura, K., Matsuoka, H. and Ando, K. (1994) Changes in haemolymph protein and their synthetic activities during development and engorgement, and after Juvenile hormone treatment in the kissing bug, *Rhodnius prolixus*. *J. Insect Physiol* 40(6): 461–499.
- Cotton, G. and Anstee, J. H. (1991) A biochemical and structural study on the effects of methoprene on fat body development in *Locusta migratoria*. *J. Insect Physiol* 37(7): 525–539.

- Elekonich, M. M., JeZ, K., Ross, A. J. and Robinson, G. E. (2002) Larval JH treatment affects pre-adult development but not adult age at onset of foraging in worker honey bees (*Apis mellifera*). *Peptides* **23**(11): 1981–90.
- Ershova, L. S., Serzhanov, O. S., Kogerman, A. P., Zolotava, S. I. and Tkachenko, A. F. (1989) The action of a synthetic JHa on the flea *Xenopsylla skjabini*. *Med Parazitol (Mosk)* **5**: 85–7.
- Gilbert, L. I., Granger, N. A. and Roe, R. M. (2000) The Juvenile hormones: historical facts and speculation on future research directions. *Insect Biochem and Mol. Biol.* **30**(8-9): 617–644.
- Ismail, S. M., Goin, C., Muthumani, K., Kim, M., Dahm, K. H. and Bhaskaran, G. (2000) Juvenile hormone acid and ecdysteroid together induce competence for metamorphosis of the Verson's gland in *Manduca sexta*. *J. Insect Physiol.* **46**(1): 59–68.
- Ismail, S. M., Satyanarayana, K., Bradfield, J. Y., Dahm, K. H. and Bhaskaran, G. (1998) Juvenile hormone acid: evidence for a hormonal function in induction of vitellogenin in larvae of *Manduca sexta*. *Arch Insect Biochem Physiol* **37**(4): 305–14.
- Jones, D. (1985) The endocrine basis for developmentally stationary prepupae in larvae of *Trichoplusia ni* pseudoparasitized by *chelonius insularis*. *J. Comp. Physiol.* **155**: 235–240.
- Kumaran, A. K. (1990) Modes of action of juvenile hormone at cellular and molecular levels. In: *Morphogenetic Hormones of Arthropods*, Gupta, A. P. (Ed). Rutgers Univ. Press: New Jersey, vol **I**: 197–198.
- Laemmli, U. K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature, Lond.* **22**: 680–685.
- Laufer, H. and Borst, W. (1983) *Juvenile Hormone and its mechanism of action. Endocrinology of Insects*, Alan R. Liss, Inc., Fifth avenue: New York, 203–216.
- Olmstead, A. W. and LeBlane, G. A. (2001) Temporal and quantitative changes in sexual reproductive cycling of the *Daphnia magna* by a juvenile hormone analog. *J. Exp. Zool.* **290**(2): 148–155.
- Short, J. E. and Edwards, J. P. (1992b) Effect of hydroprene on development and reproduction in the Oriental Cockroach, *Blatta Orientalis*. *Med Vet Entomol* **6**(3): 244–50.
- Tiwari, L. D. (1989) Effect of Diflubenzuron (Dimilin) treatment on the haemolymph proteins of *Diacrisia oblique* (walker). *Indian J. Ent.* **51**: 163–166.
- White, P. F. and Czajikowska, M. (2001) Effects of Methoprene and Permethrin on paedogenetic larvae of *Heteropza pygmaea* and *Mycophila speyeri* (Diptera: cecidomyiidae). *J. Econ. Entomol.* **93**(40): 1222–1226.
- Wyatt, G. R. and Davey, K. G. (1996) Cellular and molecular actions of JH II. Roles of JH in adult insects. *Adv. In. Insect. Physiol.* **26**: 1–155.

(Received 14 May 2007; accepted 7 September 2007)



Assessment of storage losses in wheat due to stored grain insect pests in Himachal Pradesh

K. R. S. Thakur, P. K. Mehta and Meena Thakur*

*Department of Entomology, CSK Himachal Pradesh Krishi Vishvavidyalaya,
Palampur, Himachal Pradesh 176 062, India
Email: mthakur51@yahoo.com*

ABSTRACT:

A survey conducted in Western Himachal Pradesh during 2001–2002 revealed that different storage structures used by farmers could be ranked in effectiveness as metal bin, metal drum with lid, metal drum without lid, wooden boxes and gunny bags based on percentage damage of grains and per cent loss in weight. Maximum number of damaged grains was found during rainy season and it was followed by summer and winter seasons. Similar trend was observed in per cent weight loss of wheat grains also. © 2007 Association for Advancement of Entomology

KEYWORDS: grain damage, weight loss in stored wheat

The damage caused by insects to stored food grains in tropical and temperate regions of the world is very common. Dry and cold climatic conditions have been reported to be unfavourable for majority of the stored grains pests in stores. Himachal Pradesh offers a favourable climate for storage insects. Grains are still stored by the farmers in traditional storage structures (Doharey, 1994). Agarwal *et al.* (1981) carried out a survey of the storage losses in Himachal Pradesh and since then no information is available regarding storage conditions and losses in the state. The present study was undertaken to fulfil this lacuna.

Survey for the present study was carried during 2001–2002 in two districts *viz.*, Kangra and Mandi in the state. Ninety samples were collected from each district following statistical norms. Per cent grain damage and percentage loss in weight were calculated. Per cent grain damage was calculated as per formula given by Adams and Schulten (1978). Per cent weight loss was assessed following the method of Harris and Lindbland (1978).

As evident from table 1 per cent damage to wheat grains due to insect pests varied from 2.98–13.91 and 3.06–17.90 in Kangra and Mandi district, respectively. Mean per cent damage was more during rainy season *i.e.* 10.53, 10.59 which was significantly

*Corresponding address: Department of Entomology and Apiculture, Dr. Y.S. Parmar University of Horticulture and Forestry Nauni, Solan 173 230, H.P., India

TABLE 1. Effect of different storage structures on the percentage of wheat grain damage and weight loss due to stored grain pests

Storage structures	Kangra						Mandi											
	Percentage of grain damage			Percentage of weight loss			Percentage of grain damage			Percentage of weight loss								
	Winter	Summer	Rainy	Mean	Winter	Summer	Winter	Summer	Rainy	Mean	Winter	Summer						
Metal bin	2.98 (1.73)	5.45 (2.32)	7.35 (2.71)	5.26 (2.25)	0.93 (0.96)	1.47 (1.21)	2.36 (1.54)	1.58 (1.24)	3.06 (1.75)	5.53 (2.35)	6.68 (2.58)	5.09 (2.23)	0.94 (0.97)	1.42 (1.18)	2.47 (1.57)	1.61 (1.24)		
Metal drum (with lid)	3.33 (1.82)	7.22 (2.69)	8.17 (2.86)	6.24 (2.46)	1.02 (1.01)	1.95 (1.39)	2.82 (1.68)	1.93 (1.36)	3.26 (1.80)	6.11 (2.47)	7.69 (2.73)	5.69 (2.34)	1.04 (1.02)	2.25 (1.49)	2.69 (1.64)	1.98 (1.39)		
Metal drum (without lid)	3.77 (1.94)	7.44 (2.73)	9.19 (3.03)	6.80 (2.57)	1.71 (1.31)	2.59 (1.60)	3.28 (1.81)	2.52 (1.57)	3.65 (1.91)	7.29 (2.69)	9.74 (3.02)	6.89 (2.54)	1.73 (1.31)	3.11 (1.76)	3.38 (1.84)	2.74 (1.64)		
Wooden box	4.11 (2.05)	8.45 (2.90)	10.99 (3.31)	7.85 (2.76)	2.14 (1.46)	3.16 (1.78)	4.40 (2.09)	3.28 (1.78)	4.11 (2.03)	9.35 (3.05)	10.94 (3.23)	8.13 (2.77)	2.05 (1.43)	3.69 (1.92)	4.42 (2.09)	3.37 (1.82)		
Gunny bag	7.15 (2.67)	13.91 (3.72)	6.97 (4.12)	12.67 (3.51)	3.05 (1.75)	5.88 (2.42)	8.15 (2.86)	5.69 (2.34)	7.10 (2.66)	15.52 (3.92)	17.90 (4.32)	13.51 (3.63)	2.87 (1.69)	5.75 (2.39)	8.08 (2.84)	5.56 (2.31)		
Mean	4.26 (2.04)	8.49 (2.87)	10.53 (3.21)		1.77 (1.30)	3.01 (1.68)	4.20 (2.00)		4.24 (2.03)	8.76 (2.90)	10.59 (3.18)		1.73 (1.29)	3.24 (1.75)	4.20 (2.00)			
CD's	St. St. 0.016 Seasons: 0.016			St. St. 0.022 seasons: 0.017			St. St. 0.087 Seasons: 0.068			St. St. 0.017 Seasons: 0.013								

different from summer 8.49, 8.76 and winter 4.26, 4.24 % at Kangra and Mandi district, respectively. Similar was the case for per cent loss in weight which ranged from 0.93–8.15 and 0.94–8.08 at Kangra and Mandi district, respectively. Maximum loss in weight was in rainy season i.e. 4.20 at both the districts, followed by summer and winter season. Thus from the data it is evident that both the per cent damage and per cent loss in weight were more in rainy season in both the districts. Per cent damage up to 13.91 and 17.90 per cent due to insect attack has been recorded in both the districts in the present study. Bains *et al.* (1976) reported 9.02–14.48 per cent damage to stored wheat grains in Punjab whereas Srivastava *et al.* (1973) have reported up to 30.1 per cent damage to kernels due to insect attack in U.P. In the present survey per cent loss in weight in the range of 0.93–8.15 was recorded in both the districts, maximum mean percent weight loss being in rainy season (4.20%) which is in conformity with Bains *et al.* (1976), Agarwal *et al.* (1981) and Singh *et al.* (1994) in adjoining states. Girish *et al.* (1975) reported 0.06–9.7 per cent grain weight loss in stored wheat in U.P.

The storage structures differed significantly in per cent damage and per cent weight loss due to insects. Per cent mean damage was minimum in metal bin followed by metal drum with lid, metal drum without lid, wooden box and gunny bag i.e. 5.26, 6.24, 6.80, 7.85, 12.67 and 5.09, 5.69, 6.89, 8.13, 13.51 per cent at Kangra and Mandi district, respectively. Similar was the case for mean per cent weight loss which was maximum in gunny bag 5.69, 5.56 per cent and minimum in metal bin 1.58, 1.61 per cent at Kangra and Mandi district, respectively. Thus the metallic structures (metal bin and metal drum with lid) are safe for storage of food grains than the traditional structures. Present findings get support from the observations of Bains *et al.* (1976) and Dhaliwal (1977) who also recorded more damage and loss in weight in traditional local storage structures as compared to gunny and wooden boxes, which may be due to their permeability to atmospheric air and moisture which enhances insect multiplication. Metal bins and metal drums with lid are tight structures and deficient in moisture. Singh *et al.* (1994) also observed that grain losses were more pronounced in traditional local containers compared to modern storage structures. They also confirmed minimum grain weight loss in metal bins as is the case with the present findings. The dependence of storage losses on the type of storage containers used has also been confirmed beyond doubt by Prakash *et al.* (1987).

REFERENCES

- Adams, J. M. and Schulten, G. G. M. (1978) Losses caused by insects, mites and microorganisms. In: *Post Harvest Grains Assessment Methods*, American Association of Cereal Chemists: 193.
- Agarwal, R. K., Singh, K. N. and Srivastva, P. K. (1981) Studies on the assessment of storage losses in wheat due to stored grain insect pests in Himachal Pradesh. *Bulletin of Grain Technology* **19**(3): 198–200.
- Bains, S. S., Battu, G. S. and Atwal, A. S. (1976) Distribution of *Trogoderma granarium* evets and other stored grain insect pests in Punjab and losses caused by them. *Bulletin of Grain Technology* **14**(1): 18–29.

- Dhaliwal, G. S. (1977) Incidence of storage insect pests in rural areas. *Indian Journal of Entomology* **39**(2): 114–116.
- Doharey, R. B. (1994) Assessment of storage losses in food grains in Punjab, Haryana and Himachal Pradesh. *Bulletin of Grain Technology* **32** (1): 3–18.
- Girish, G. K., Jain, S. K., Kumar, A. and Aggarwal, N. S. (1975) Part V: Assessment of storage losses, quality and pesticidal contamination in wheat available in the markets of Western Uttar Pradesh, Punjab and Haryana. *Bulletin of Grain Technology* **32** (1): 30–31.
- Harris, K. L. and Lindbland, C. O. (1978) *Post Harvest Grain Loss Assessment Methods*, American Association of Cereal Chemist: Washington D.C., U.S.A., 193.
- Prakash, A., Rao, J., Pasalu, I. C. and Mathur, K. C. (1987) *Rice Storage and Insect Pests Management*, B.R. Publishing Corporation: New Delhi, 328.
- Singh, G., Singh, J., Thopar, V. K., Sehgal, V. K. and Paul, S. (1994) Storage and losses of wheat grains at farm level in Punjab. *Bulletin of Grain Technology* **32** (1): 18–19.
- Srivastava, P. K., Tripathi, B. P., Girish, G. K. and Krishnamurthy, K. (1973) Studies on the assessment of losses III. Conventional grain storage practices and losses in rural areas in Western Uttar Pradesh. *Bulletin of Grain Technology* **11**(2): 129–139.

(Received 18 January 2007; accepted 19 May 2007)



The suitability of host plants for mass rearing of *Scirtothrips dorsalis* Hood (Thysanoptera: Thripidae)

P. N. Ganga Visalakshy* and A. Krishnamoorthy

*Division of Entomology and Nematology, Indian Institute of Horticultural Research,
Hessarghatta Lake post, Bangalore 560 089, India
Email: gangesv@iihar.ernet.in*

ABSTRACT: A mass rearing method for *Scirtothrips dorsalis* on rose flower is described. French bean pods and brinjal leaves were found unsuitable for rearing.

© 2007 Association for Advancement of Entomology

KEYWORDS: *Scirtothrips dorsalis*, mass rearing

Scirtothrips dorsalis Hood (Thysanoptera: Thripidae) is a serious pest of chilli, brinjal, rose and other economically important crops in India (Ananthakrishnan, 1971; Jaganmohan *et al.*, 1980). Life stages desap tender leaves, shoots, developing flowers and fruits causing leaf curl upwards symptom (Amin, 1976), resulting in 46–90% yield loss in sweet and chilli pepper (Krishnakumar, 1995). This insect is also a key vector of tomato spotted wilt virus (TSWV) which causes bud necrosis disease (Amin *et al.*, 1981). The present study was carried out to develop a mass rearing method for *S. dorsalis*.

Scirtothrips dorsalis was obtained as follows: Terminal shoots about 10 cm long of field grown chilli plants, showing typical leaf curl symptom were cut and placed in a tall cylindrical glass jar. The mouth of the jar was closed with a muslin cloth to prevent escape of adults. The container was placed in a B.O.D incubator, at 10 °C for 10 min. to inactivate the insects. The twig was then taken out and tapped gently over a clean black platform. Most of the thrips fell down to the platform. Adults alone were collected in a clean tall glass jar (15 × 10 cm). From this sample, *S. dorsalis* were isolated based on the taxonomic characters and used for mass rearing.

Three host plants, brinjal leaves, French bean pods and rose were selected for the study. These are the natural host plants of *S. dorsalis* (Ananthakrishnan, 1971). The host plants were grown in glasshouse and parts of plants, free from any insect, including thrips, were used for the experiments.

*Corresponding author

Rearing on French bean pods: Tender, washed and air dried pods were spread on top of a brass wire mesh placed on top of a filter paper, kept moist with wet cotton inside a plastic jar. *S. dorsalis* females were released and after two days adults were recovered and released into new rearing jars. Bean pods were kept undisturbed for emergence of larvae. Fresh bean pods were added to the jar as and when the exposed ones dried. Observations on the number of larvae, pupae and adults produced were recorded. The experiment was replicated five times with two adults per rearing jar.

Rearing on brinjal leaves: Fresh tender brinjal leaf was placed on the concave side of a watch glass flanked with a blotting paper. The edge of the leaf was trimmed, so that the leaf remained within the watch glass. This was later kept inside a circular petri plate (20 cm diameter \times 0.5 cm height) containing water. The cut end of the stalk was covered with a cotton wick that was in contact with water in the petri dish. After releasing *S. dorsalis* females on leaves, the petri plate was covered with another one of the same size. The released adults were recovered after two days and the process was repeated. Fresh leaf was kept as and when the previous one started drying. Observations on the number of larvae, pupae and adults at each stage were recorded. The experiment was replicated five times with five adults per replication.

Rearing on rose: Flowers of var. Happiness, Mulu and Gladiator were chosen. Half opened flowers which were free from any insect attack were selected. A plastic vial with lid (5 \times 3 cm) was taken and 3/4th of its capacity was filled with water. A circular hole was made in the centre of the lid, through which the flower stalk was inserted. A petri plate (20 width \times 0.5 cm height) was taken and a watch glass was kept centrally with the concave surface facing upwards. The plate was filled with water up to half its capacity. The plastic vial with the flower was placed on top of the watch glass. The flower was covered with a glass chimney (10 \times 15 cm width/height). After releasing *S. dorsalis* the other end was covered with a tissue paper held tight by rubber band.

To record the progeny production, the exposed flower was taken out at weekly intervals and tapped gently on to a watch glass (15 cm diameter) that was kept inside a petri plate (25 \times 0.5 w/h) filled with water. Majority of thrips fall to the watch glass. Tapping was continued for two or three times till no more thrips fell down. The flower was later replaced into the rearing chamber and observation was repeated after a week by following the same procedure.

The number of larvae and pupae falling to the watch glass was recorded. Larvae were released into fresh rearing chambers for further development, while pupae were kept on a rose petal inside a Petri-plate (10 cm diam.) having a moist filter paper, folded into two or three layers, for adult emergence. The experiment was carried out with two females per replication and each treatment was replicated six times.

Among the three host plants tried, french bean pod and brinjal leaves were found to be unsuitable for multiplication of *S. dorsalis*. In french bean pods, though released females were observed to remain alive up to a week, no larvae could be recorded indicating absence of oviposition. No larvae were recorded in brinjal leaf also. The insects multiplied on rose flower. Among the rose varieties, Gladiator and Mulu were

TABLE 1. Progeny production and per cent survival of *S. dorsalis* at different generations

Generations	No of larvae recorded	No. of pupae recorded	No of adults recorded
1st	98.40	87.60	73.40
2nd	83.00	73.80	67.20
3rd	85.40	87.20	61.40
5th	85.60	73.20	61.00
10th	81.20	73.90	59.00
CV	10.69	14.76	22.17
CD @ 0.05%	13.02	15.66	19.14

found to be equally suitable. Happiness flowers wilted within two or three days and were not suitable for mass rearing.

Preliminary studies on variety Gladiator, showed that releasing two females per flower yielded more progeny (60 nymphs/female) as compared to releasing five (29 nymphs/flower) or ten females per flower (16 nymphs/female). Hence further studies were carried out by releasing two females per flower.

Progeny production in different generations are presented in table 1. The data showed no deterioration in progeny production up to 10th generation.

Observations on rose flowers showed that larval stages were more in inner petals than in outer petals. Pre-pupae and pupae occurred on the undersurface of sepals, on curled tips of petals and on the base of petals. Normally pre-pupae and pupae were found aggregated together while larval stages were dispersed on the flower.

As already noted among the host plants studies, rose var. Gladiator was the best for rearing of *S. dorsalis*. Regular changing of flower was not needed as a single flower could support all stages of *S. dorsalis* for more than 10 days, without wilting. As and when the exposed flower starts drying fresh flower was kept on top of the earlier one. Thrips migrated within 1–2 h, to the new flower. After 24 h the old flower was removed and culture was continued on the fresh one.

ACKNOWLEDGEMENTS

The authors are grateful to the Director, Indian Institute of Horticultural Research, Bangalore, for the facilities and encouragement provided during the course of the study and to Dr. Vikas Kumar, Research Associate, Division of Entomology and Nematology, for identification of *S. dorsalis*.

REFERENCES

- Amin, P. W. (1976) Leaf curl disease of chilli peppers in Maharashtra, India. *Pests articles and News summaries* 25: 131–134.

- Amin, P. W., Reddy, D. V. R., Ghanekar, A. M. and Reddy, M. S. (1981) Transmission of tomato spotted wilt virus, causal agent of bud necrosis of peanut by *Scirtothrips dorsalis* and *Frankliniella schultzei*. *Plant Disease* **65**: 663–665.
- Ananthakrishnan, T. N. (1971) Thrips (Thysanoptera) in agriculture, horticulture and forestry—diagnosis, bionomics and control. *Journal of Scientific and Industrial Research* **30**: 113–140.
- Jaganmohan, N., Krishnaiah, K. and Prasad, V. G. (1980) Evaluation of insecticides for the control of pests of sweet pepper (*Capsicum annum* var. *Grosum* Linn). *Pestology* **5**: 1–4.
- Krishnakumar, N. K. (1995) Yield loss in chilli and sweet pepper due to *Scirtothrips dorsalis* Hood (Thysanoptera: Thripidae). *Pest Management in Horticultural Ecosystems* **1**: 61–70.

(Received 31 May 2006; accepted 22 February 2007)



Ant fauna associated with whiteflies in some locales of Eastern and Western Ghats in two Southern states of India

R. W. Alexander Jesudasan and M. Anto Claver*

Department of Zoology, Madras Christian College, Chennai 600059, India

ABSTRACT: A list of whiteflies attracting ant species on wide range of flora in some localities of Eastern and Western Ghats of Tamil Nadu and Kerala is provided. The ecotypic diversity of ant tending whiteflies such as *Aleuromarginatus* sp., *Aleuroclava* sp., *Sphercaleyrodes indica*, and *Rhachisphora* sp. on the host plants was influenced by species of tending ants. The whitefly, *Aleuroclava* sp. inviting ant species were totally different in the same host plant of distant localities whereas the whitefly *Aleuromarginatus* sp. attracting ant species in the nearby localities were similar. The whitefly *Rhachisphora* sp. attracted diverse ant species from different localities.

© 2007 Association for Advancement of Entomology

KEYWORDS: mutualistic interaction, ant tending whitefly, Western Ghats

Understanding the interactions between species or populations is a prerequisite for predicting ecological phenomena at all levels of biological organization. For this respect, mutualistic interactions have generally received less attention as compared to competition and predation in the concept of species existence and survival among living organisms. In most mutualisms, one partner performs some service that benefits its associate, and in return receives some pay-off or reward. The local biotic environment, that is, the identity and abundance of other species with which the mutualists interact, also strongly influences outcomes. Costs and benefits of anti-herbivore mutualisms are influenced by the herbivores' host plant (Stadler *et al.*, 2001).

Hemipterans infesting a wide range of host plants exude honeydew, which attract a number of ants belonging to the Subfamilies Dolichoderinae, Formicinae, Myrmicinae and Pseudomyrmicinae (Veeresh and Ali, 1990; Rastogi *et al.*, 1997). With some aleurodids species viz., *Rhachisphora*, *Sphercaleyrodes*, *Aleuromarginatus*, *Aleuroclava*, *Aleurolobus* and *Dialeurodes*, ants are found frequently associated for honeydew and wax (Jesudasan and David, 1995). Whiteflies benefit from ant tending because the presence of ants deter insect predators from their host plants, increase their feeding

*Present address: Department of Zoology, St. Andrew's College, Gorakhpur 273001, India

TABLE 1. Whiteflies attracting ant species in some localities of Eastern and Western Ghats

Location/Host plants	Whitefly	Ants and their subfamilies
Aliyar/ <i>Grewia</i> sp.	<i>Aleuromarginatus</i> sp.	Formicinae <i>Anoplolepis gracilipes</i> Jerdon <i>Camponotus compressus</i> Fabr. <i>Camponotus rufoglaucus</i> Jerdon Myrmicinae <i>Crematogaster</i> sp. Ponerinae <i>Pachycondyla</i> sp. Pseudomyrmicinae <i>Tetraponera nigra</i> Jerd
Top Slip/ <i>Grewia</i> sp	<i>Aleuromarginatus</i> sp.	Formicinae <i>Anoplolepis gracilipes</i> Jerdon <i>Camponotus compressus</i> Fabr. <i>Camponotus rufoglaucus</i> Jerdon Myrmicinae <i>Crematogaster</i> sp. Ponerinae <i>Pachycondyla</i> sp.
Mannuthu/ <i>Cassia</i> sp.	<i>Aleuroclava</i> sp.	Dolichoderinae <i>Tapinoma melanocephalum</i> Fabr. Formicinae <i>Camponotus compressus</i> Fabr. Myrmicinae <i>Crematogaster rothneyi</i> Mayr <i>Monomorium indicum</i> Fornel <i>M. latinode</i> Mayr Pseudomyrmicinae <i>Tetraponera rufonigra</i> Jerdon
Edapalyam/ <i>Cassia</i> sp.	<i>Aleuroclava</i> sp.	Formicinae <i>Paratrechina longicornis</i> Latr. <i>Monomorium</i> sp.
Mekkarai/ <i>Ochlandra</i> <i>travancorica</i>	<i>Sphercaleyrodes indica</i>	Dolichoderinae <i>Techomyrmex albiceps</i> Smith Formicinae <i>Acantholepis</i> sp. <i>Anoplolepis gracilipes</i> Jerdon <i>Anoplolepis longipes</i> Fabr. <i>Paratrechina</i> sp. <i>Polyrhachis illudata</i> Walker <i>P. rastallata</i> Latr <i>P. exercita</i> Walker Myrmicinae <i>Catulacus taprobannea</i> Smith

TABLE 1. (Contd...)

Location/Host plants	Whitefly	Ants and their subfamilies
Aryankavu/ <i>Ochlandra travancorica</i>	<i>Sphericalleyrodes indica</i>	Formicinae <i>Anoplolepis gracilipes</i> Jerdon <i>Anoplolepis longipes</i> Fabr. <i>Paratrechina longicornis</i> Latr. <i>Polyrhachis</i> sp. <i>Monomorium</i> sp.
Pamalakovu/ <i>Ochlandra travancorica</i>	<i>Sphericalleyrodes indica</i>	Dolichoderinae <i>Techomyrmex albiceps</i> Smith Formicinae <i>Anoplolepis longipes</i> Jerdon <i>Polyrhachis rastallata</i> Latr. <i>Oecophylla smaragdina</i> Fabr. Myrmicinae <i>Myrmecaria brunnea</i> Saunders Pseudomyrmicinae <i>Tetraponera aitkeni</i> Forel
Damdam Parai/ <i>Ixora pavetta</i>	<i>Rhachisphora</i> sp.	Dolichoderinae <i>Techomyrmex</i> sp. <i>Tapinoma melanocephalum</i> Fabr. Formicinae <i>Camponotus compressus</i> Fabr. Myrmicinae <i>Crematogaster</i> sp. <i>Myrmecaria brunnea</i> Saunders
Megamalai <i>I. pavetta</i> , <i>Memecylon umbellatum</i> , <i>Syzygium cumini</i>	<i>Rhachisphora</i> sp.	Dolichoderinae <i>Tapinoma melanocephalum</i> Fabr. Formicinae <i>Camponotus compressus</i> Fabr. Myrmicinae <i>Crematogaster rothneyi</i> Mayr. <i>Monomorium indicum</i> Forel <i>M. latinode</i> Mayr Pseudomyrmicinae <i>Tetraponera rufonigra</i> Jerdon

rates, and prevent the buildup of fungus from collected honeydew (Delabie, 2001). However, the nature of interaction can exhibit considerable spatial variation (Cushman *et al.*, 1998), and it is therefore important to observe this interaction across such special scale. Hence, a study was undertaken to collect and identify the whiteflies and ants from some host plants occurring in different localities of Eastern Ghats and Western Ghats of Tamil Nadu and Kerala, India.

TABLE 1. (Contd...)

Location/Host plants	Whitefly	Ants and their subfamilies
Tambaram	<i>Rhachisphora</i> sp.	Dolichoderinae
<i>I. pavetta</i> ,		<i>Tapinoma melanocephalum</i> Fabr.
<i>M. umbellatum</i> ,		Formicinae
<i>Syzygium cumini</i>		<i>Camponotus paria</i> Emery
		<i>Camponotus rufoglaucus</i> Jerdon
		<i>Camponotus</i> sp. nr. <i>taylori</i>
		<i>Oecophylla smaragdina</i> Fabr.
		<i>Paratrechina longicornis</i> (Latr.)
		<i>Polyrhachis exercita</i> (Walker)
		<i>P. scissa</i> Roger
		Myrmicinae
		<i>Crematogaster dalyi</i> Forel
		<i>Meranoplus bicolor</i> Guer
		<i>Monomorium indicus</i> Forel
		<i>Myrmicaria brunnea</i> Saunders.
		<i>Pheidole</i> sp.
		Pseudomyrmicinae
		<i>Tetraponera aitkeni</i> (Forel)
		<i>Tetraponera nigra</i> Jerdon
		<i>Tetraponera rufonigra</i> (Jerdon)

Collection of leaves bearing whitefly tending ants were made from Tambaram (Se-laiyur range), Damdam Parai (Palani Hills), Aryankavu and Edapalayam (Thenmalai range), Mekkarai (Tenkasi range), Mannuthu (Megamalai range), Aliyar (Valparai range), Top slip (Anaimalai Hills), Pamalakavlu (Adimali range), on the host plants viz., *Ixora pavetta*, *Memecylon umbellatum* (Burman), *Syzygium jambolanum* (Lam.), *S. cumini*, *Grewia* sp., *Cassia* sp. and the reed, *Ochlandra travancorica* (Bedd.).

The ecotypic diversity of ant tending whiteflies such as *Aleuromarginatus* sp., *Aleuroclava* sp., *Sphercaleyrodes ochlandrae*, and *Rhachisphora* sp. on the host plants was influenced by species of tending ants (Table 1). The ant, *Anoplolepis longipes* was commonly tending the whitefly host plants on all the three localities whereas *Catulacus taprobannea*, *Acantholepis* were only found in Mekkarai. The ant *Techomyrmex albiceps* was tending the whitefly, *Sphercaleyrodes indica* in Mekkarai and Pamalakavlu localities whereas *Oecophylla smaragdina*, *Myrmicaria brunnea* and *Tetraponera aitkeni* were only seen in Pamalakavlu local whitefly species. The ants such as *Polyrhachis* sp. and *Paratrechina* sp. were tending the whitefly of Mekkarai and Aryankavu where as *Monomorium* sp. was only noticed in Aryankavu site host plant. Intraspecific variation in host-plant quality as influenced by environmental conditions might affect herbivore-ant interactions (Pierce *et al.*, 1991).

The ants attending to the whitefly, *Aleuroclava* sp. were totally different on same host plants from distant localities such as Mannuthu (Magamalai) and Edapalayam (Thenmalai) whereas the whitefly *Aleuromarginatus* sp. attracting ant species in the

nearby localities viz., Aliyar (Valparai), Top slip (Anaimalai) of the host plant, *Grewia* sp., were almost similar (Table 1). Cushman *et al.* (1998) stated that some members of homopteran species were significantly more to attract ants than those of others.

Diverse ants have been noted to be associated with the whitefly, *Rhachispora* sp. from different localities (Table 1). The *Tapinoma* sp., *Crematogaster* sp. and *Camponotus* sp. were common in occurrence in all the localities (Damdam Parai, Mannuthu and Tambaram). Similarly, Cushman and Addicott (1991) reported that homopteran attracting ants were often varied in their assemblages and taxonomic diversity across large geographic spatial scale.

ACKNOWLEDGEMENTS

The assistance rendered by Dr. B. Ravichandran and Dr. A. Jayasanker are acknowledged. We thank the Ministry of Environment and Forests, New Delhi for the financial support through a research project (23/41/2001-RE). Thanks to Dr. T. M. Musthak Ali, University of Agricultural Sciences, Bangalore for the identification of ants and Dr. D. Narasimhan, Department of Botany, MCC for identifying the flora.

REFERENCES

- Cushman, J. H., Compton, S. G., Zachariades, C., Ware, A. B., Nefdt, R. J. C. and Rashbrook, V. K. (1998) Geographic and taxonomic distribution of a positive interaction: attended homopterans indirectly benefit figs across southern Africa. *Oecologia* **116**: 373–380.
- Cushman, J. H. and Addicott, J. F. (1991) Conditional interactions in ant-plant-herbivore mutualisms. In: *Ant-Plant Interactions*, Huxley, C. R. and Cutler, D. F. (Eds). Oxford Univ. Press: 92–103.
- Delabie, J. H. C. (2001) Trophobiosis between Formicidae and Hemiptera (Sternonophyncha and Auchenorrhyncha): an overview. *Neotropic entomology* **30**(4): 501–516.
- Jesudasan, R. W. A. and David, B. V. (1995) Association of aleyrodids with aleyrodids and other insects. *Journal of Applied Zoological Research* **6**(1): 49–50.
- Pierce, V. E., Nash, D. R., Baylis, M. and Carper, E. R. (1991) Variation in the attractiveness of lycaenid butterfly larvae to ants. In: *Ant-Plant Interactions*, Huxley, C. R. and Cutler, D. F. (Eds). Oxford Univ. Press: 131–142.
- Rastogi, N., Nair, P., Kolatkar, M., Willa, H. and Godagkar, R. (1997) Ant fauna of the Indian Institute of Science Campus—Survey and some preliminary observation. *Journal of Indian Institute of Science* **77**: 133–140.
- Stadler, B., Dixon, A. F. C. and Kindlmann, P. (2001) Relative fitness of aphids: effects of plant quality and ants. *Ecology Letters* **5**: 216–222.
- Veeresh, G. K. and Ali, T. M. (1990) The present status of our knowledge on Indian ants. In: *Social Insects: An Indian Perspective*, Veeresh, G. K., Kumar, A. R. V. and Shivashankar, T. (Eds). IUSSI—Indian Chapter, Bangalore, India: 77–102.

(Received 28 August 2006; accepted 6 March 2007)



Histochemical study of yolk granules of ripe oocytes of *Oxya hyla hyla* (Orthoptera: Acrididae)

K. Das and D. Ghosh*

Insect Physiology and Biochemistry Laboratory, Department of Life Sciences, Tripura University, Suryamaninagar 799130, Tripura, India
Email: ghoshdurgadas@rediffmail.com

ABSTRACT: Isolated yolk granules of vitellogenic oocyte of *Oxya hyla hyla* under light microscope revealed the presence of deeply stained proteinaceous rim region and lightly stained core region. The same granules were also positive for carbohydrate and lipid. Those yolk granules initially deposited in smaller size gradually showed increase in size due to aggregation of different components.

© 2007 Association for Advancement of Entomology

KEYWORDS: *Oxya hyla hyla*, ripe oocyte, yolk granules, histochemistry

Vitellogenesis is one of the key events occurring in growing oocytes of both vertebrates and invertebrates. This process is usually under control of several hormones (Courceles and Kondo, 1980). Although a good number of works have been published on purification and characterization of yolk proteins (vitellin, lipovitellin) (Harnish and White, 1982) including that of Locust (Chen *et al.*, 1978) sufficient attention has not been paid to study the yolk granules taken from the ripe oocyte in its original condition under microscope. Raikhel (1984) using high-resolution immuno- and cytochemical study had shown that in mosquito (*Aedes aegypti*) oocytes the yolk granules are initially of low electron density but gradually grow larger in size and of higher electron density. *Oxya hyla hyla* (grasshopper) is an important orthopteran paddy pest of Tripura. Depending on structure and physiology, the ovariole of *Oxya* is panoistic in nature and is without any nurse cells. Ghosh *et al.* (1997) by histological studies had shown that in growing oocytes of *Gesonula punctifrons*, the small sized yolk droplets initially deposited near the follicular epithelium slowly grew larger in size. Therefore an effort was made to study the isolated yolk granules under light microscope using various cytochemical methods.

Ripe terminal oocytes were collected from the mature females of *Oxya hyla hyla* collected from paddy fields around Agartala city. For histochemical studies, yolk granules were obtained by rupturing the fully ripe oocytes over a slide and

*Corresponding author




FIGURE 1. Photomicrograph of sectioned vitellogenic oocyte of *Oxya hyla hyla* stained with Hg-Bromphenol Blue showing the yolk granules (Yg) of different sizes. 350X.

FIGURE 2. Photomicrograph of the separated yolk granules (Yg) stained with Hg-Bromphenol Blue showing differential stainability of the peripheral rim (pr) and lightly staining core (c). 330X.

FIGURE 3. Photomicrograph of the deeply stained Sudan Black B positive granules (Ld) depicting the lipoidal nature of the yolk granules. 330X.

FIGURE 4. Photomicrograph showing the PAS positive nature of the overall granule depicting presence of carbohydrate substance in the yolk granules. 390X.

spreading its contents homogeneously over it. Then those yolk contents which were seen to contain spheres in unstained condition under microscope were fixed in Ca-formol solution and air dried for 24 h or more. Those slides were used for further histochemical studies. To examine the presence of proteins in the yolk granules lipids were extracted from this slide by methanol-chloroform (1:1) extraction method. Protein moiety in the granules was stained by Hg-Bromphenol blue method (Pearse, 1975). To stain lipid granules, Sudan black B was used. In this staining method, three sets of Ca-formol fixed dried slides containing the yolk granules (one set treated with methanol-chloroform extraction method, one set treated with cold acetone extraction method and one set without any extraction) were first immersed in 2.5% aqueous Bromine solution for 30 min and then washed in distilled water for 30 min. Then

the slides were immersed in 70% alcohol for 30 min and stained in saturated Sudan Black B solution in 70% alcohol. Excess stain was removed by washing in distilled water (Pearse, 1975). To stain mucoid substances in the yolk granules Schiff's reagent was prepared and staining was done after oxidation by Periodic acid solution (Pearse, 1975).

The yolk granules when stained with Hg-Bromphenol blue showed intensely Hg-Bromphenol blue positive reaction. The diameter of those granules were about $32.8 \pm 9.3 \mu\text{m}$. It was also observed that those Hg-Bromphenol blue positive granules showed differential staining reaction in its various zones (Fig. 1, 2). The inner core region of those granules became faintly stained while the outer rim became deep blue. The diameter of the core region was about $22 \pm 5 \mu\text{m}$ and the outer rim was about $5.5 \pm 2 \mu\text{m}$ thick. This result indicated that the granules were composed of different types of protein components. The basic protein originally becomes more intensely stained in comparison to acidic or neutral proteins (Mazia *et al.*, 1953). So it was apparent that the outer rim of those protein granules were made up of proteins with more basicity and in the inner core region basicity was low. Again when those yolk granules were stained with Sudan Black B the granules treated with methanol-chloroform and cold acetone extraction methods showed negative reaction but the set without any extraction showed strong positive reaction and became brownish black in colour (Fig. 3). This result suggested the presence of lipid or lipoprotein in the yolk granules. Further when stained by PAS reaction after oxidation with Periodic acid those granules of almost same diameter showed PAS positive reaction and became faint magenta in colour (Fig. 4). Such positive reaction suggested that those yolk granules also contained carbohydrate moiety.

All the results taken together strongly suggested that the yolk granules or the droplets which were present in the oocyte had a complex nature and constituted of protein, carbohydrate and lipid moieties. In the sectioned condition (Fig. 1) these granules were found in different sizes and in isolated condition also, that size variation was apparent. The variation of the size of the yolk granules strongly suggested that there was a morphogenetic pathway in the development of the large size yolk granules from the smaller. That may be due to aggregation of different components or other phenomena may be involved. Raikhel (1984) using high resolution immuno and cytochemical electron microscopy have shown, a similar pathway in mosquito oocyte. He has also indicated that in the yolk granules some crystalline and some noncrystalline protein part was present. In the present situation, the presence of proteinaceous core and the rim suggested the same phenomena.

ACKNOWLEDGEMENT

The authors are thankful to the North Eastern Council for providing a Scholarship to KD and also to the authorities of the Tripura University for providing the laboratory facilities.

REFERENCES

- Anderson, E. (1964) Oocyte differentiation and vitellogenesis in the roach *Periplaneta americana*. *The Journal of Cell Biology* **20**(1): 131–155.
- Chen, T. T., Strahlendorf, P. W. and Wyatt, G. R. (1978) Vitellin and vitellogenin from Locusts (*Locusta migratoria*). *The Journal of Biological Chemistry* **253**(15): 5325–5331.
- Courceles, D. F. and Kondo, M. (1980) Lipovitellin of the crustacean, *Artemia salina*: Biochemical analysis of Lipovitellin complex from the yolk granules. *The Journal of Biological Chemistry* **255**(14): 6727–6733.
- Dejmal, R. K. and Brookes, V. J. (1972) Insect Lipovitellin. Chemical and physical characteristics of yolk protein from the ovaries of *Leucophaea maderae*. *The Journal of Biological Chemistry* **247**(3): 860–874.
- Ghosh, D., Chel, G. and Pal, S. G. (1997) Histophysiology of ovarian development and oogenesis in *Gesonula puctifrons* (Orthoptera:Acrididae). *Uttar Pradesh Journal of Zoology* **17**(2): 139–145.
- Harnish, D. G. and White, B. N. (1982) Insect vitellin: Identification, purification and characterization from eight orders. *Journal of Experimental Zoology* **220**: 1–10.
- Mazia, D., Brewer, P. A. and Alfert, M. (1953) The cytochemical staining and measurement of protein with Hg-Bromphenol Blue. *Biol. Bull. Mar. Boil. Lab., Woods hole* **104**: 57–67.
- Pearse, A. G. E. (1975) *Histochemistry*, 3rd edition. Churchill Livingstone: Edindurge London and New York, 647–648.
- Raikhel, A. S. (1984) The accumulative pathway of vitellogenin in the Mosquito oocyte: A High-resolution Immuno- and cytochemical study. *Journal of Ultrastructure Research* **87**: 285–302.

(Received 20 March 2007; accepted 5 June 2007)

GUIDELINES TO AUTHORS

Scope: ENTOMON will publish original research papers on insects, arachnids and myriapods. Reviews are not acceptable.

Papers on morphology, anatomy and histology will be considered only when they form part of systematics, physiology or behavioural studies.

Announcements of seminars/symposia, book reviews and other items of entomological interest will also be considered for publication.

Types of papers: Articles up to 3 printed pages in length (1800 words) will be published as **short communications** and those between 4 and 10 pages (up to 6000 words) as **full papers**.

Full paper should be organized under the following sub-heads—Abstract, Introduction, Materials and Methods, Results, Discussion, Acknowledgements and References.

Short communication should be organized in the same way, but without the sub-headings.

Publication policy: Manuscripts submitted for publication in ENTOMON should not have been published or submitted for publication elsewhere.

At least one of the authors should be a member of AAE.

Page charges are applicable at the rate of Rs. 100 per printed page for authors in India and US \$ 10 for authors overseas. Invoice will be sent to the authors along with the galley proof.

Twenty five **reprints** of the paper will be supplied free of cost to the authors.

Typing and submission: Manuscripts should be typed double space on one side of good quality paper, having 3.5 cm left margin and 2.5 cm right margin.

The first page should contain the title, author's name affiliation and e-mail address. When the number of authors is more than one, indicate the name of the corresponding author with an asterisk and specify 'Corresponding author' in a footnote. The second page should contain the abstract, followed by key words and a running title. From page 3 onwards, type the text continuously from Introduction to References. Place the tables on separate sheets at the end of the paper. The pages should be numbered.

Three copies of the manuscript, complete in all respects including illustrations, should be sent to the Managing Editor, ENTOMON, CABB, University of Kerala Kariavattom, P.O., Thiruvananthapuram 695 581, Kerala, India.

An electronic version need be sent only after acceptance of the paper for publication.

Guide for writing: Careful attention to the following guide will facilitate early acceptance of your paper for publication in ENTOMON. Although some of these suggestions may appear trivial, they have been prompted by our experience in reviewing the papers received for publication. Keep in mind that ENTOMON is a research journal and your paper will be read only by those who are specialists in the respective fields of research.

Title should be brief and should reflect the specific content of the research reported in the paper.

Abstract should be informative, not indicative. It should very briefly highlight the aim of the study and major conclusions. Normally it should not exceed 150 words.

Key words should be limited to four or five most pertinent indicators of the work, relevant to indexing the article.

Introduction should be brief, normally not exceeding 300 words. Include the specific aim of the research, a review of the available information in the area and existing gap in knowledge. The introduction should thus justify the work carried out. Avoid elementary details and repetition of well known facts. For example, in a paper reporting the efficacy of a biopesticide against a particular pest, it is not necessary to explain the hazards of chemical pesticides, alternative methods of insect control and advantage of integrated pest management. Limit the literature review to what is most relevant to the topic under study.

Materials and Methods should provide just enough details to permit proper interpretation of the results. Materials used need not be described separately if it is evident from the methods given. Technical description of the method is needed only when the method is new. If the method followed has been already described elsewhere, just give the reference. If any alteration is made, describe the alteration alone, with reason.

Results: Adequate care should be taken to organize and present the results in a clear, concise and summarized form.

Quantitative data should always be analysed using suitable statistical methods. Organize the data into well planned tables. Each table should be self-explanatory.

Do not repeat the data presented in the table in the text. Quote the relevant figures in the text only when it is essential for highlighting some particular finding.

Due care should be taken while interpreting the results of statistical analysis. For example, treatments which show higher numerical value cannot be treated as superior to those having lower numerical values when there is no statistically significant difference.

Interpretation of the data should be with reference to the objectives set in the experiment.

Do not include graphs duplicating the data presented in the tables.

When the research involves repetition of the work already reported by others, include the new findings alone in the paper.

Illustrations included in the paper should be essential for explaining some points in the text. Photographs of the life stages of an insect are not useful unless the insect is being reported for the first time. Illustration should be of good quality. Limit the number of photographs to 4–6 and organize them into plates wherever possible. The illustrations should be numbered consecutively as Fig. 1, Fig. 2, etc., without distinction between drawings, graphs and photographs. Labelling should be legible and large enough to stand suitable reduction. Legend for the figures should be typed on a separate page. All figures must be referred to, at appropriate places, in the text.

The cost of printing colour illustration is to be met by the author.

Discussion: The discussion section is intended to critically analyse and interpret the results with reference to the objectives set forth in the study. It should highlight the importance of the results in relation to what is already known. It should also point out the limitations of the study, if any. The discussion should not repeat details given under Results, except to highlight some conclusions.

References should list all publications cited in the text, in alphabetical order. Do not include any references not cited in the text. Some authors show a tendency to cite too many references in support of a statement. Cite only a few references most pertinent to the point dealt with. Follow the citation style given below.

Examples of citations in text:

Krishnaswamy (1978, 1979)

Govindan *et al.* (1998)

(Reddy 1978; David 1991)

Examples of citations under References:

Articles in Journals:

Nayar K. K. (1953) Neurosecretion in *Iphita*. *Current Science* 22(2): 149.

Nair M. R. G. K. and Mohandas N. (1962) On the biology and control of *Carvalhoeia arecae*, a pest of areca palms in Kerala. *Indian Journal of Entomology* 24: (1) 86–93.

Jalaja M. Muraleedharan D. and Prabhu V. K. K. (1973) Effect of extirpation of median neurosecretory cells on reproduction in the female red cotton bug, *Dysdercus cingulatus*. *Journal of Insect Physiology* 19(1): 29–36

Books and Articles in Books:

Novak V. J. A. (1966) *Insect Hormones*. Methuen and Co., 478 pp.

Wigglesworth V. B. (1964) The hormonal regulation of growth and reproduction in insects. In: *Advances in Insect Physiology* Vol. 2 (Eds. Beament J. W. L., Treherne J. E. and Wigglesworth V. B.), Academic Press, London, pp 247–335.

Review Procedure: Manuscripts will be subjected to a preliminary scrutiny by an Editorial team and those conforming to the general guidelines set to above will be referred to experts for detailed review; others will be returned to the author for resubmission in the prescribed format. Based on detailed review, the corresponding author will be advised of, acceptance, non-acceptance or need for revision. The revised

manuscript will be scrutinized again by an Editorial team (and by expert referees, if needed) before final acceptance. On final acceptance, the author will be asked to submit an electronic version of the manuscript. Proof will be sent to the corresponding author. It should be checked and returned within 3 days of receipt. The journal reserves the right to proceed with publication if corrections are not communicated promptly.

Strict Conformity with the above Guidelines will Ensure Speedy Publication of Your Paper.

ENTOMON

ENTOMON is a quarterly journal of the Association for Advancement of Entomology issued in March, June, September and December, devoted to publication of research work on various aspects of insects and other arthropods.

EDITORIAL ADVISORY BOARD

T. N. ANANTHAKRISHNAN, Emeritus Scientist, Chennai
G. BHASKARAN, A & M University, Texas
K. P. GOPINATHAN, Indian Institute of Science, Bangalore
ZUO. R. SHEN, Agricultural University, Beijing

EDITORIAL BOARD

HOFFMANN, K. H., Germany
SEIJI TANAKA, Japan
ABRAHAM, C. C., Thrissur
APARNA DUTTA GUPTA, Hyderabad
DUBEY, O. P., Delhi
GADAGKAR, R., Bangalore
MOHANDAS, N., Trivandrum
NARENDRA, T. C., Calicut
NAIR, K. S. S., Trivandrum
PALANISWAMI, M. S., Trivandrum
PRABHU, V. K. K., Trivandrum
RABINDRA R. J., Bangalore
RAMAMURTHY, V. V., Delhi
SASIDHARAN PILLAI, K., Trivandrum
MURALEEDHARAN, D. (Managing Editor)
MARIAMMA JACOB (Associate Editor)

Address MS and all editorial correspondence to Managing Editor, ENTOMON, Centre for Arthropod Bioresources and Biotechnology (CABB), University of Kerala, Kariavattom, Trivandrum 695581, India.

SUBSCRIPTION RATES

Annual subscription for Institutions: Rs. 2000.00 (in India); US\$ 250 (Air Mail)
Annual subscription for individuals: Rs. 300.00 (in India); US\$ 100 (Air Mail)

© 2007 by the Association for Advancement of Entomology. All rights reserved

1. All remittance to the Journal or Association should be sent to the Secretary-Treasurer of the Association by Bank Draft only, A/c payee in favour of the Association for Advancement of Entomology, payable at Kariavattom.
2. Requests for replacement copies of ENTOMON in lieu of numbers lost in transit, should reach the Secretary-Treasurer not later than three months after the date of publication of the number.

ENTOMON is covered in the following abstracting/indexing journals: *Chemical Abstracts*, *Review of Applied Entomology*, *Science Citation Index* and *Current Contents/Agriculture, Biology and Environmental Sciences*, *Biological Abstracts*, *Entomology Abstracts* and other relevant abstracts, *Referativny Zhurnal* and *Current Advance in Biological Sciences* and *New Entomological Taxa*.

Reg. No. 123SBN
